

SLAC-PUB-5886  
LBL-32725  
LU TP 92-22  
DTP 92-58  
August 1992  
(T/E)

MLLA and the Average Charged Multiplicity of  
Events Containing Heavy Quarks in  $e^+e^-$  Annihilation\*

Bruce A. Schumm

*Physics Division*

*Lawrence Berkeley Laboratory, Berkeley, California 94720*

Yuri L. Dokshitzer\*

*Department of Theoretical Physics*

*University of Lund, Sölvegatan 14-A, S-22362 Lund, Sweden*

Valery A. Khoze \*

*Center For Particle Theory*

*University of Durham, Durham DH1 3LE, U.K.*

and

*Stanford Linear Accelerator Center, Stanford, California, 94309*

Dale S. Koetke

*CERN, European Organization for Particle Physics, CH-1211 Geneva 23, Switzerland*

Submitted to *Physical Review Letters*

\* Work supported in part by the Department of Energy contracts DE-AC03-76SF00098 (LBL), and DE-AC03-76SF00515 (SLAC), and the SERC of the United Kingdom.

\* On leave of absence from the Institute for Nuclear Physics, St. Petersburg, Gatchina 188350, Russia

## ABSTRACT

We present the MLLA results for heavy quark event multiplicities in  $e^+e^-$  annihilation, which provide that the difference  $\delta_{Ql}$  between heavy and light quark event multiplicities at the same cms energy should be a calculable constant, independent of cms energy. Published data on heavy quark event multiplicities are presented in this light, and are consistent with the energy independence of the multiplicity difference. Averaging over all cms energies, we find that  $\delta_{bl} = 4.3 \pm 0.9$  and  $\delta_{cl} = 2.2 \pm 1.2$  tracks, while the corresponding MLLA expectations are  $5.5 \pm 0.8$  and  $1.7 \pm 0.5$  tracks, respectively. We briefly discuss the experimental prospects for improving this test of MLLA, and suggest further experimental tests of MLLA-LPHD modelling of heavy quark events.

The perturbative (PT) approach to QCD jet physics is based on the Modified Leading Logarithmic Approximation (MLLA), which incorporates both double- and single-logarithmic effects in the development of parton cascades.<sup>[1-3]</sup> In addition, the hypothesis of Local Parton-Hadron Duality (LPHD),<sup>[1][4]</sup> which is supported by experimental studies of multi-hadron production in QCD jets,<sup>[5]</sup> suggests a close correspondence between the observable inclusive characteristics of hadron spectra and those calculated at the parton level by means of PT QCD. Thus, when combined with LPHD, MLLA can hope to describe the gross features of hadronic systems, such as multiplicity distributions, the angular distribution of particle flows, inclusive energy spectra, etc., without invoking phenomenological fragmentation schemes. In this approach, nonperturbative effects are reduced to normalizing coefficients relating hadronic characteristics to partonic ones, which, according to LPHD, must be independent of both the hardness of the initiating partons and the energy range of the final state particles.

Until now the main phenomenological successes of the MLLA-LPHD approach were connected with the description of the inclusive characteristics of jets in  $e^+e^-$  annihilation, without distinction between the contributions of light and heavy primary quarks.<sup>[6,7]</sup> Prompted by these successes, and the recent availability of data on heavy quark jets, one would like to compare the PT predictions for heavy quark generated jets with existing data on heavy quark events. In this Letter, then, we shall present a comparison between the MLLA-LPHD description of particle multiplicity with data on the mean charged multiplicity of events containing heavy hadrons.

The physics of heavy quarks has always been considered a particularly good laboratory for detailed studies of QCD. The large quark mass  $M_Q \gg \Lambda_{QCD}$  provides a natural cut-off, which keeps the relevant space-time region compact enough to avoid the truly strong, non-PT domain of strong interactions. In the case of  $e^+e^-$  annihilation at center of mass (cms) energies  $W \gg M_Q \gg \Lambda_{QCD}$ , one can hope for a good description of many inclusive properties of hadronic jets via MLLA-LPHD.

The results of the PT description of specific properties of particle distributions in heavy quark jets have been announced in a number of publications,<sup>[6][8-10]</sup> but only recently has this subject been addressed more comprehensively.<sup>[11]</sup> It was demonstrated<sup>[6][9][11]</sup> that the difference in many properties of hadronic jets produced by heavy quarks (excluding the products of the weak decay of the heavy quark itself), from that of light (u,d,s) quarks, originates from the restriction of the phase space available to gluon radiation associated with the kinematic effects of the heavy quark mass.

Particle multiplicities were calculated<sup>[6][11]</sup> by convoluting the differential distribution of gluon ( $g$ ) radiation from a massive quark ( $Q$ ), with energy  $E_Q \gg M_Q$ , with the MLLA-LPHD multiplicity distribution of subsequent gluon cascades. The angular radiation pattern of soft gluons, with energy  $\omega \ll E_Q$  and emission angle  $\Theta \ll 1$ ,

$$d\sigma_{Q \rightarrow Q+g} \sim \frac{\Theta^2 d\Theta^2}{[\Theta^2 + \Theta_0^2]^2} \frac{d\omega}{\omega} \quad (1)$$

where

$$\Theta_0 = \frac{M_Q}{E_Q}, \quad (2)$$

gives rise to the familiar double logarithmic approximation (DLA)<sup>[3]</sup> for  $\Theta > \Theta_0$ :

$$d\sigma_{Q \rightarrow Q+g} \sim \frac{d\Theta^2}{\Theta^2} \frac{d\omega}{\omega} = d(\ln \Theta^2) d(\ln \omega). \quad (3)$$

For  $\Theta < \Theta_0$ , however, the angular integration is no longer logarithmic, and the yield of particles in this region from (1) adds only a small ( $O(\sqrt{\alpha_s} \cdot N)$ ) contribution to the total multiplicity  $N$ . This region of suppressed radiation in the forward direction is known as the ‘dead cone’.<sup>[6]</sup> On the other hand, for emission angles  $\Theta \gg \Theta_0$ , and for the internal structure of secondary gluon jets (due to the strict angle-ordering of hadronic cascades), Equation (1) yields completely identical behavior between light and heavy quark gluon radiation. This universality of the gluon radiation spectrum, up to a depopulation in a cone around  $Q$  of opening angle  $\Theta \sim \Theta_0$ ,

lies at the heart of comparisons between light and heavy quark jets within the framework of MLLA.

A consequence of this suppression of forward gluon radiation is that the ‘companion’ multiplicity  $\Delta N(Q\bar{Q}; W)$  of light hadrons accompanying the heavy quark, excluding the decay products of the on-shell heavy hadron, is less than the particle yield in a light quark jet at the same cms energy. Quantitatively, to MLLA accuracy,<sup>[6][11]</sup>

$$\Delta N(Q\bar{Q}; W) = N(q\bar{q}; W) - N(q\bar{q}; \sqrt{\epsilon} \cdot M_Q) + O(\alpha_s(M_Q^2)N(q\bar{q}; M_Q)) \quad (4)$$

where  $N(q\bar{q}; W)$  is the mean total multiplicity in light quark events at cms energy  $W$ , and  $\epsilon = \exp(1)$ . It should be noted that this MLLA calculation improves the DLA calculation of Ref. 9, which was limited in accuracy by higher order corrections of  $O(\sqrt{\alpha_s(W^2)}N(q\bar{q}; W))$ . The uncertainty from this correction is substantially larger than the  $O(\alpha_s(M_Q^2)N(q\bar{q}; M_Q))$  leading correction of the MLLA approach, and is formally a function of cms energy.

The most important consequence of this MLLA result stems from the fact that both the subtrahend and uncalculated higher-order correction in Equation (4) are functions of the fixed mass scale  $M_Q$ , and thus independent of the cms energy  $W$  of the  $e^+e^-$  annihilation. Thus, it is a fundamental prediction of MLLA that the difference

$$N(q\bar{q}; W) - \Delta N(Q\bar{Q}; W) \quad (5)$$

is almost completely independent of the cms energy  $W$ ; leading  $W$ -dependent corrections are of  $O(\sqrt{\alpha_s(W^2)} \cdot M_Q^2/W^2)$ .<sup>[12]</sup> In addition, the extraction of  $N(q\bar{q}; \sqrt{\epsilon} \cdot M_Q)$  from existing low cms energy multiplicity data permits an estimate of the mean multiplicity difference (5) to  $O(\alpha_s(M_Q^2)N(q\bar{q}; M_Q))$  accuracy.

Viewed another way, it is QCD coherence, which consideration of the gluonic formation length shows to apply to the region  $\Theta \lesssim \Theta_0 \equiv M_Q/E$ , that provides for this relation between light and heavy quark multiplicities. The difference between

light and heavy quark radiation in this forward region, where gluons radiated from heavy quarks can not distinguish themselves quantum-mechanically from the heavy parent quark, is roughly the integral of the light quark radiation spectrum out to  $\Theta_0$ , which is dominated by the  $N(q\bar{q}; \sqrt{e} \cdot M_Q)$  term in Equation (4).

It should be emphasized that, due to the observed steeply rising dependence of total multiplicity on  $\ln(W)$ , the MLLA multiplicity picture is *not* consistent with the naively expected reduction of the energy scale<sup>[8][13]</sup>

$$\Delta N(Q\bar{Q}; W) = N(q\bar{q}; (1 - \langle x_Q \rangle)W) \quad (6)$$

where  $\langle x_Q \rangle = 2\langle E_Q \rangle/W$ . A good description of the total multiplicity data is provided by the MLLA inspired multiplicity formula<sup>[1-3]</sup>

$$N(q\bar{q}; W) \propto \alpha_s(W^2)^{\frac{1}{4} + \frac{10n_f}{27b}} \cdot \exp\left(\frac{\sqrt{96\pi}}{b} \cdot \alpha_s^{-\frac{1}{2}}(W^2)\right) \quad (7)$$

where  $n_f$  is the number of quark flavors and  $b = 11 - 2n_f/3$ . Applying this to Equation (6), the difference (5) is not constant, but is asymptotically proportional to  $N(W)$ :

$$N(q\bar{q}; W) - \Delta N(Q\bar{Q}; W) \simeq \sqrt{\frac{6\alpha_s(W^2)}{\pi}} \cdot \ln \frac{1}{1 - \langle x_Q \rangle} \cdot N(q\bar{q}; W). \quad (8)$$

With the recent addition to previous results<sup>[14-17]</sup> of a measurement at the  $Z^0$  resonance by the MARK II collaboration,<sup>[18]</sup> the average total charged multiplicity  $\bar{n}_b$  in  $e^+e^-$  annihilation to  $b$  quarks has been measured in a range of cms energy from  $W = 29 \text{ GeV}$  to  $W = 91 \text{ GeV}$ . In addition, the average total charged multiplicity  $\bar{n}_c$  for  $e^+e^- \rightarrow c\bar{c}$  events has been measured by two experiments at  $W = 29 \text{ GeV}$ .<sup>[15][16]</sup> Combined with the world sample of total hadronic (all quark flavors)  $e^+e^-$  mean charged multiplicity measurements ( $\bar{n}_{had}$ ) between  $W = 1.5 \text{ GeV}$  and  $W = 91 \text{ GeV}$ ,<sup>[19]</sup> these data can be used to study the difference in charged particle yields between light (u,d,s) and heavy (c,b) quark production.

Table 1 shows the measured mean charged multiplicity of events containing heavy quarks at  $W = 29, 35, 42.1, \text{ and } 90.9 \text{ GeV}$ . Also shown is the total hadronic multiplicity at the same energies, derived from the corresponding multiplicity data in the cms energy region surrounding the heavy quark multiplicity point. Both heavy quark-associated and total hadronic multiplicities have been corrected for the effects of initial state radiation (ISR), so that the quoted values correspond to the average charged multiplicity that would be observed at the given cms energy in the absence of ISR. Charged tracks from  $K_s^0$  and  $\Lambda$  decays are included in the measured multiplicities.

The difference  $\delta_{Ql} \equiv \bar{n}_Q - \bar{n}_l$  between measured heavy and light quark event total charged multiplicities can be written as

$$\delta_{Ql} = \frac{f_l \cdot \bar{n}_Q + f_b \cdot \bar{n}_b + f_c \cdot \bar{n}_c - \bar{n}_{had}}{f_l} \quad (9)$$

where  $Q = c, b$ , and the  $f_i$  are the standard model production fractions for light, c, and b quarks. For  $\gamma^*$  decays,  $f_l = 0.55$ ,  $f_c = 0.36$ ,  $f_b = 0.09$ , while at the  $Z^0$ ,  $f_l = 0.61$ ,  $f_c = 0.17$ ,  $f_b = 0.22$ .

In order to calculate  $\delta_{bl}$  from the measurements of  $\bar{n}_b$  and  $\bar{n}_{had}$ , it is necessary to estimate  $\bar{n}_c$  in the unmeasured region above  $W = 29 \text{ GeV}$ . For this purpose, we have made the MLLA-motivated assumption that  $\bar{n}_c - \bar{n}_l$  is constant as a function of cms energy, which gives  $\bar{n}_c = 14.7 \pm 0.7$  and  $16.0 \pm 0.7$  tracks at  $W = 35$  and  $42.1 \text{ GeV}$ , respectively. At these energies, the uncertainty in  $\bar{n}_c$  is dominated by the  $\pm 0.7$  track uncertainty in  $\bar{n}_c$  at  $W = 29 \text{ GeV}$ . Due to the large difference in cms energy between  $90.9$  and  $29 \text{ GeV}$ , in order to maintain model independence we have assumed only that  $\bar{n}_l < \bar{n}_c < \bar{n}_b$ , leading to a value  $\bar{n}_c = 22.1 \pm 2.5$  tracks at  $W = 90.9 \text{ GeV}$ . Because of the relatively small contribution of  $Z^0 \rightarrow c\bar{c}$  (.17 of  $\sigma_{had}$ ), and the large statistical error in  $\bar{n}_b$  at  $90.9 \text{ GeV}$ , this constitutes only a small contribution to the uncertainty in  $\delta_{bl}$ .

Combining these values for  $\bar{n}_c$  with the  $\bar{n}_{had}$ ,  $\bar{n}_c$  and  $\bar{n}_b$  results in Table 1 yields the results for  $\delta_{cl}$  and  $\delta_{bl}$  exhibited in Table 2. Also shown are averages for

all experiments at  $W = 29 \text{ GeV}$ , and for all cms energies combined, taking into account common systematic errors due to the uncertainty in the average  $\bar{n}_c$ , ISR corrections, the common use of lepton tagging to identify heavy quark events at  $W = 29 \text{ GeV}$ , and the common use of displaced vertex information to identify  $b$  quark events at  $W = 35 \text{ GeV}$  and  $W = 42.1 \text{ GeV}$ . The results for the individual measurements of  $\delta_{bl}$  are also displayed in Figure 1. To the available accuracy, the results are seen to be independent of energy, in marked contrast to the steeply rising total multiplicity data, and are thus consistent with the MLLA prediction discussed above. On the other hand, Figure 1 in Ref. 18 exhibits a mild (1.1 standard deviation) disagreement between the energy dependence of the  $Z \rightarrow b\bar{b}$  companion multiplicity and that of the naive expectation of Equation (6).

Since Equation (4) is a statement about the *companion* multiplicity of events containing heavy quarks, it should be noted that any energy dependence of the heavy hadron decay multiplicity will modify the claim that  $\delta_{Ql}$  is a constant. Since the heavy quark-associated multiplicity results discussed here were measured at cms energies well above the heavy hadron thresholds, such effects are not expected to be large.

Equation (4) predicts that the difference between the total light quark and companion heavy quark event multiplicities should be equal to the total light quark event multiplicity at  $W = \sqrt{e} \cdot M_Q$ . In terms of the heavy hadron decay multiplicity  $\bar{n}_Q^{dk}$ , this can be written

$$\delta_{Ql} \equiv \bar{n}_Q - \bar{n}_l = \bar{n}_Q^{dk} - \bar{n}_l(\sqrt{e} \cdot M_Q). \quad (10)$$

In order to estimate  $\bar{n}_l(\sqrt{e} \cdot M_Q)$ , we assume  $\sqrt{e} \cdot M_c = \sqrt{e} \cdot 1.5 = 2.5 \text{ GeV}$  and  $\sqrt{e} \cdot M_b = \sqrt{e} \cdot 4.8 = 7.9 \text{ GeV}$ , and use the measured total hadronic multiplicity for  $2 < W < 3 \text{ GeV}$ <sup>[20,21]</sup> and  $5.5 < W < 10 \text{ GeV}$ <sup>[21,22]</sup> respectively, to estimate  $\bar{n}_{had}$  at these cms energies. At  $W = 2.5 \text{ GeV}$ , below the charm threshold,  $\bar{n}_{had} = \bar{n}_l$ , while for  $W = 7.9 \text{ GeV}$ , we assume the value of  $\delta_{cl}$  measured at  $W = 29 \text{ GeV}$  to



correct  $\bar{n}_{had}$  for the effects of the 40% admixture of  $c\bar{c}$  events. This yields

$$\bar{n}_l(\sqrt{e} \cdot M_c) = 3.5 \pm 0.4 \text{ tracks} \quad (11)$$

$$\bar{n}_l(\sqrt{e} \cdot M_b) = 5.5 \pm 0.7 \text{ tracks.} \quad (12)$$

For  $W = \sqrt{e} \cdot M_b$ , the quoted uncertainty includes a 0.5 track contribution from the uncertainty in the  $c\bar{c}$  correction.

Assuming the values  $\bar{n}_c^{dk} = 5.2 \pm 0.3$ <sup>[23]</sup> and  $\bar{n}_b^{dk} = 11.0 \pm 0.2$ <sup>[18]</sup> yields

$$\bar{n}_c^{dk} - \bar{n}_l(\sqrt{e} \cdot M_c) = 1.7 \pm 0.5 \text{ tracks} \quad (13)$$

$$\bar{n}_b^{dk} - \bar{n}_l(\sqrt{e} \cdot M_b) = 5.5 \pm 0.8 \text{ tracks.} \quad (14)$$

It should be noted that  $O(\alpha_s \cdot \bar{n}_l(M_Q))$  terms neglected in Equation (4) are expected to be of roughly the same size as the experimental uncertainties on these values. Comparing these values to the results for  $\delta_{cl}$  and  $\delta_{bl}$  in Table 2, it is again seen that the experimental data are consistent with the predictions of MLLA.

In conclusion, it has been seen that, to within the available accuracy, the observed mean multiplicities of events containing heavy hadrons are in good agreement with the predictions of MLLA, and in mild disagreement with the naive relation (6). In particular, the data support the notion that the difference between the companion multiplicity in heavy quark events, and the total multiplicity in light quark events at the same cms energy, is independent of cms energy. This provides a fundamental check of the consistency of the MLLA approach, which predicts this result and provides that it should be independent of higher order corrections. In addition, combined with the quantitative agreement between this multiplicity difference and the lower cms energy multiplicity data embodied in Equation (10), this work supports the validity of LPHD as a phenomenological approach to modelling confinement.

Based on the result from Ref. 18, which was statistically limited, experiments currently running at 91 GeV at LEP and the SLC should be able to measure  $\bar{n}_b$  to  $\pm 0.5$  tracks or better. Combined with a measurement of  $\bar{n}_c$  to  $\pm 1.0$  tracks, this would yield a measurement of  $\delta_{bl}$  to  $\sim \pm 0.8$  tracks, providing a much more stringent test of the MLLA predictions in the case of  $b$  quark production. Further reduction of the uncertainty in  $\bar{n}_c$  to  $\pm 0.5$  tracks or better would allow measurements of  $\delta_{cl}$  and  $\delta_{bc}$  to  $\sim \pm 0.7$  tracks, providing stringent tests of MLLA-LPHD predictions down to the  $M_c^2$  scale. At this lower mass, the question of the relationship between LPHD and QCD confinement becomes particularly interesting.

Making use of the inclusive properties of heavy hadron decay to statistically remove the heavy hadron decay tracks, it should be possible to study more extensively the properties of radiated hadrons in heavy hadron events. It is expected that the gluonic radiation 'dead cone' will appear as a depopulation in the region  $x_Q > \Lambda_{QCD}/M_Q$ , while the spectrum of soft hadrons with  $x_Q \ll \Lambda_{QCD}/M_Q$  should be identical to that of light quark jets.<sup>[6][10][11]</sup> Finally, MLLA-LPHD predict various aspects of the  $x_Q$  distribution itself,<sup>[24]</sup> which can in principle be tested with an accurate measurement of  $\langle x_Q \rangle$  and the  $x_Q$  spectrum.

In closing, we would like to express thanks to Michael Peskin and John Jaros of SLAC for interesting discussions and helpful comments. One of us (V.K.) would like to thank the SLAC Theory Group for hospitality during his visit. This work was supported in part by U.S. Department of Energy contracts DE-AC03-76SF00098 (LBL), and DE-AC03-76SF00515 (SLAC), and the SERC of the United Kingdom.

## REFERENCES

1. Yu. L. Dokshitzer and S.I. Troyan, Proc. XIX LNPI Winter School, V.I. (1984) 144; preprint LNPI-922 (1984).
2. A. H. Mueller, Nucl. Phys. **B213** (1983) 85, and corrections, Nucl. Phys. **B241** (1984) 141.
3. Yu. L. Dokshitzer, V.A. Khoze, A.H. Mueller, and S.I. Troyan, *Basics of Perturbative QCD*, ed. Tran Than Van, Editions Frontieres, Paris, 1991.
4. Ya. I. Azimov *et al.*, Z. Phys. **C27** (1985) 65; Z. Phys. **C31** (1986) 213.
5. For recent reviews, see J. Drees, G. Cowan, talks at the workshop *QCD - 20 Years Later*, June 9-13, 1992, Aachen.
6. Yu. L. Dokshitzer, V.A. Khoze, and S.I. Troyan, J. Phys. G: Nucl. Part. Phys. **17** (1991) 1481; *ibid.*, p. 1602.
7. Yu. L. Dokshitzer, V.A. Khoze, and S.I. Troyan, Z. Phys **C55** (1992), 107.
8. Ya. I. Azimov, Yu. L. Dokshitzer, and V. A. Khoze, Sov. J. Nucl. Phys. **36** (1982) 878; Ya. I. Azimov *et al.*, *ibid.* **37** (1983) 419.
9. Yu. L. Dokshitzer, V. S. Fadin, and V. A. Khoze, Phys. Lett. **B115** (1982) 242; *ibid.*, Zeit. Phys. **C15** (1982) 325.
10. Yu. L. Dokshitzer, V. A. Khoze, S. I. Troyan, in: *Proceedings of the 6<sup>th</sup> Intl. Conf. on Physics in Collision*, p. 417, ed. M. Derrick, World Scientific, Singapore, 1987.
11. Yu. L. Dokshitzer, V. A. Khoze, S. I. Troyan, Specific Features of Heavy Quark Production: I. Leading Quarks, LU TP 92-10; II. LPHD Approach to Heavy Particle Spectra, LU TP 91-13 (under preparation); III. Light Hadron Accompaniment, LU TP 91-16 (under preparation).
12. See Item III in Ref. 11.
13. A. V. Kisselev *et al.*, Z. Phys. **C41** (1988), 521.

14. M. Sakuda *et al.*, Phys. Lett. **B152** (1985), 399.
15. P. C. Rowson *et al.*, Phys. Rev. Lett. **54** (1985), 2580.
16. H. Aihara *et al.*, Phys. Lett. **B134** (1987), 299.
17. W. Braunschweig *et al.*, Z. Phys. **C42** (1989), 17.
18. B. A. Schumm and D. S. Koetke *et al.*, Phys. Rev. **D46** (1992), 453.
19. For a compilation of total multiplicity data, see Ref. 18.
20. C. Bacci *et al.*, Phys. Lett. **B86** (1979), 234.
21. J. L. Siegrist *et al.*, Phys. Rev. **D26** (1982), 969.
22. B. Niczyporuk *et al.*, Z. Phys. **C9** (1981), 1.
23. The value used is that of Ref. 16, with an addition of 0.1 tracks to account for the effects of higher mass charm states.
24. See Item II in Ref. 11.

## Table Captions

Table 1: Measured mean charged multiplicities. For the total multiplicity  $\bar{n}_{had}$ , the value is from an average of all experiments in the cms energy region surrounding the heavy quark event multiplicity point.

Table 2: Derived differences between heavy and light quark event mean multiplicities.

EXPERIMENT	$E_{cm}$ (GeV)	$\bar{n}_{had}$	$\bar{n}_c$	$\bar{n}_b$
DELCO <sup>[14]</sup>	29.0	$12.41 \pm 0.21$		$14.3 \pm 1.2$
MARK II <sup>[15]</sup>	29.0	$12.41 \pm 0.21$	$13.2 \pm 1.0$	$16.1 \pm 1.1$
TPC <sup>[16]</sup>	29.0	$12.41 \pm 0.21$	$13.5 \pm 0.9$	$16.7 \pm 1.0$
TASSO <sup>[17]</sup>	35.0	$13.59 \pm 0.30$		$16.0 \pm 1.5$
TASSO <sup>[17]</sup>	42.1	$14.85 \pm 0.40$		$17.0 \pm 2.0$
MARK II <sup>[18]</sup>	90.9	$20.94 \pm 0.20$		$23.1 \pm 1.9$


TABLE 1

EXPERIMENT	$E_{cm}$ (GeV)	$\delta_{cl}$	$\delta_{bl}$
DELCO <sup>[14]</sup>	29.0		$2.9 \pm 1.5$
MARK II <sup>[15]</sup>	29.0	$1.9 \pm 1.7$	$5.0 \pm 1.4$
TPC <sup>[16]</sup>	29.0	$2.4 \pm 1.5$	$5.7 \pm 1.3$
AVERAGE	29.0	$2.2 \pm 1.2$	$4.7 \pm 1.0$
TASSO <sup>[17]</sup>	35.0		$3.6 \pm 1.9$
TASSO <sup>[17]</sup>	42.1		$3.3 \pm 2.5$
MARK II <sup>[18]</sup>	90.9		$3.3 \pm 2.7$
AVERAGE	ALL ENERGIES	$2.2 \pm 1.2$	$4.3 \pm 0.9$

TABLE 2

### Figure Caption

Figure 1: Energy dependence of total multiplicity (open points) and the multiplicity difference between b and light quark production (filled points) in  $e^+e^-$  annihilation. MLLA predicts unambiguously that this multiplicity difference should be independent of energy. Also shown is the expected value of this multiplicity difference, given by lower energy multiplicity data in accordance with MLLA (see text). The one standard deviation range indicated by the dotted lines is dominated by the uncertainty in the light quark event multiplicity at  $E_{cm} = \sqrt{e} \cdot M_b$ , and does not include a  $\sim 1$  track uncertainty due to (energy independent) higher order corrections to MLLA. Citations for the total multiplicity data are compiled in Ref. 18.



XBL928-6137

Figure 1