

Strong Symmetry Breaking at e^+e^- Linear Colliders

Timothy L. Barklow*

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309 USA

(Dated: December 18, 2001)

The study of strong symmetry breaking at an e^+e^- linear collider with $\sqrt{s} = 0.5 - 1.5$ TeV is reviewed. It is shown that processes such as $e^+e^- \rightarrow \nu\bar{\nu}W^+W^-$, $e^+e^- \rightarrow \nu\bar{\nu}t\bar{t}$, and $e^+e^- \rightarrow W^+W^-$ can be used to measure chiral Lagrangian and strong resonance parameters. The linear collider results are compared with those expected from the LHC.

I. INTRODUCTION

Until a Higgs boson with large couplings to gauge boson pairs is discovered, the possibility of strong electroweak symmetry breaking must be entertained. Without such a particle the scattering of gauge bosons will become strong at a scale of order 1 TeV. The most commonly studied class of theories which deals with this scenario is technicolor [1]. A generic prediction of technicolor theories is that there is a vector resonance with mass below about 2 TeV which unitarizes the WW scattering cross section. Scalar and tensor resonances are also possible, along with light pseudo-Goldstone bosons which can be produced in pairs or in association with other particles [2].

Independent of the model, the strong interactions of gauge bosons below the threshold for resonance production can be described by an effective chiral Lagrangian in analogy with $\pi\pi$ scattering below the ρ resonance [3] :

$$\mathcal{L}_{SB} = \mathcal{L}^{(2)} + \mathcal{L}^{(4)} + \dots$$

$$\begin{aligned} \mathcal{L}^{(2)} = & \frac{v^2}{4} \text{Tr} D^\mu \Sigma^\dagger D_\mu \Sigma + \frac{g'^2 v^2}{16\pi^2} b_1 (\text{Tr} T \Sigma^\dagger D_\mu \Sigma)^2 \\ & + \frac{gg'}{16\pi^2} a_1 \text{Tr} \Sigma B^{\mu\nu} \Sigma^\dagger W_{\mu\nu} \\ \mathcal{L}^{(4)} = & \frac{\alpha_4}{16\pi^2} \text{Tr} (D_\mu \Sigma^\dagger D_\nu \Sigma) \text{Tr} (D^\mu \Sigma^\dagger D^\nu \Sigma) + \frac{\alpha_5}{16\pi^2} [\text{Tr} (D^\mu \Sigma^\dagger D_\mu \Sigma)]^2 \\ & - ig \frac{L_{9L}}{16\pi^2} \text{Tr} (W^{\mu\nu} D_\mu \Sigma D_\nu \Sigma^\dagger) - ig' \frac{L_{9R}}{16\pi^2} \text{Tr} (B^{\mu\nu} D_\mu \Sigma^\dagger D_\nu \Sigma) . \end{aligned}$$

Here $W^{\mu\nu}$ and $B^{\mu\nu}$ are related to the $SU(2) \times U(1)$ gauge fields as in [3], D_μ is the covariant derivative, g and g' are the $SU(2) \times U(1)$ coupling constants, and Σ is composed of the Goldstone boson fields w^k :

$$\Sigma = \exp \left(\frac{i w^k \tau^k}{v} \right) ,$$

where the τ^k are Pauli matrices and $v = 246$ GeV is the Standard Model Higgs vacuum expectation value parameter. The chiral Lagrangian parameters a_1 and b_1 are tightly constrained by precision electroweak data [5]. The terms with coefficients α_4 and α_5 induce anomalous quartic gauge boson couplings which can be measured by observing gauge boson scattering in processes such as $e^+e^- \rightarrow \nu\bar{\nu}W^+W^-$ and $\nu\bar{\nu}ZZ$. The terms with coefficients L_{9L} and L_{9R} induce anomalous triple gauge couplings (TGC's) which can be measured in the reaction $e^+e^- \rightarrow W^+W^-$.

In this paper we summarize strong symmetry breaking signals and the measurement of chiral Lagrangian and strong resonance parameters at an e^+e^- linear collider (LC) with a center of mass system (CMS) energy in the range of 0.5 to 1.5 TeV. Many of the results are taken from the strong symmetry breaking sections of Ref. [4], which the reader is invited to consult for further details.

*timb@slac.stanford.edu; Work supported by Department of Energy contract DE-AC03-76SF00515.

TGC	error $\times 10^{-4}$			
	$\sqrt{s} = 500$ GeV		$\sqrt{s} = 1000$ GeV	
	Re	Im	Re	Im
g_1^γ	15.5	18.9	12.8	12.5
κ_γ	3.5	9.8	1.2	4.9
λ_γ	5.4	4.1	2.0	1.4
g_1^Z	14.1	15.6	11.0	10.7
κ_Z	3.8	8.1	1.4	4.2
λ_Z	4.5	3.5	1.7	1.2

TABLE I: Expected errors for the real and imaginary parts of CP-conserving TGCs assuming $\sqrt{s} = 500$ GeV, $\mathcal{L} = 500 \text{ fb}^{-1}$ and $\sqrt{s} = 1000$ GeV, $\mathcal{L} = 1000 \text{ fb}^{-1}$. The results are for one-parameter fits in which all other TGCs are kept fixed at their SM values.

II. $e^+e^- \rightarrow \nu\bar{\nu}W^+W^-$, $\nu\bar{\nu}ZZ$, $\nu\bar{\nu}t\bar{t}$

The first step in studying strong WW scattering is to separate the scattering of a pair of longitudinally polarized W 's, denoted by $W_L W_L$, from transversely polarized W 's and background such as $e^+e^- \rightarrow e^+e^-W^+W^-$ and $e^- \bar{\nu}W^+Z$. Studies have shown that simple cuts can be used to achieve this separation in $e^+e^- \rightarrow \nu\bar{\nu}W^+W^-$, $\nu\bar{\nu}ZZ$ at $\sqrt{s} = 1000$ GeV, and that the signals are comparable to those obtained at the LHC [6, 7]. Furthermore, by analyzing the gauge boson production and decay angles it is possible to use these reactions to measure the chiral Lagrangian parameters α_4 and α_5 with an accuracy greater than that which can be achieved at the LHC [8].

The reaction $e^+e^- \rightarrow \nu\bar{\nu}t\bar{t}$ provides unique access to $W^+W^- \rightarrow t\bar{t}$ since this process is overwhelmed by the background $gg \rightarrow t\bar{t}$ at the LHC. Techniques similar to those employed to isolate $W_L W_L \rightarrow W^+W^-$, ZZ can be used to measure the enhancement in $W_L W_L \rightarrow t\bar{t}$ production [9–12]. Even in the absence of a resonance it will be possible to establish a clear signal. The ratio S/\sqrt{B} is expected to be 12 for a linear collider with $\sqrt{s} = 1$ TeV, 1000 fb^{-1} and 80%/0% electron/positron beam polarization, increasing to 22 for the same luminosity and beam polarization at $\sqrt{s} = 1.5$ TeV.

III. $e^+e^- \rightarrow W^+W^-$

Strong gauge boson interactions induce anomalous TGC's at tree-level:

$$\begin{aligned}\kappa_\gamma &= 1 + \frac{e^2}{32\pi^2 s_w^2} (L_{9L} + L_{9R}) \\ \kappa_Z &= 1 + \frac{e^2}{32\pi^2 s_w^2} \left(L_{9L} - \frac{s_w^2}{c_w^2} L_{9R} \right) \\ g_1^Z &= 1 + \frac{e^2}{32\pi^2 s_w^2} \frac{L_{9L}}{c_w^2}.\end{aligned}$$

where κ_γ , κ_Z , and g_1^Z are TGC's [13], $s_w^2 = \sin^2 \theta_w$, and $c_w^2 = \cos^2 \theta_w$. Assuming QCD values for the chiral Lagrangian parameters L_{9L} and L_{9R} , κ_γ is shifted by $\Delta\kappa_\gamma \sim -3 \times 10^{-3}$.

The TGCs can be measured by analyzing the W^+W^- production and decay angles in the process $e^+e^- \rightarrow W^+W^-$. Table I contains the estimates of the TGC precision that can be obtained at $\sqrt{s} = 500$ and 1000 GeV for the CP-conserving couplings g_1^V , κ_V , and λ_V . These estimates are derived from one-parameter fits in which all other TGC parameters are kept fixed at their tree-level SM values. For comparison the LHC with $\mathcal{L} = 300 \text{ fb}^{-1}$ is expected to measure κ_γ and κ_Z with an accuracy of 0.006 and 0.01, respectively. The 4×10^{-4} precision for the TGCs κ_γ and κ_Z at $\sqrt{s} = 500$ GeV can be interpreted as a precision of 0.26 for the chiral Lagrangian parameters L_{9L} and L_{9R} . Assuming naive dimensional analysis [14] such a measurement would provide a 8σ (5σ) signal for L_{9L} and L_{9R} if the strong symmetry breaking energy scale were 3 TeV (4 TeV).

When WW scattering becomes strong the amplitude for $e^+e^- \rightarrow W_L W_L$ develops a complex form factor F_T in analogy with the pion form factor in $e^+e^- \rightarrow \pi^+\pi^-$ [15, 16]. To evaluate the size of this effect the following

expression for F_T can be used:

$$F_T = \exp\left[\frac{1}{\pi} \int_0^\infty ds' \delta(s', M_\rho, \Gamma_\rho) \left\{ \frac{1}{s' - s - i\epsilon} - \frac{1}{s'} \right\}\right]$$

where

$$\delta(s, M_\rho, \Gamma_\rho) = \frac{1}{96\pi} \frac{s}{v^2} + \frac{3\pi}{8} \left[\tanh\left(\frac{s - M_\rho^2}{M_\rho \Gamma_\rho}\right) + 1 \right].$$

Here M_ρ, Γ_ρ are the mass and width respectively of a vector resonance in $W_L W_L$ scattering. The term

$$\delta(s) = \frac{1}{96\pi} \frac{s}{v^2}$$

is the Low Energy Theorem (LET) amplitude for $W_L W_L$ scattering at energies below a resonance. Below the resonance, the real part of F_T is proportional to $L_{9L} + L_{9R}$ and can therefore be interpreted as a TGC. The imaginary part, however, is a distinct new effect.

The real and imaginary parts of the form factor F_T are measured in $e^+e^- \rightarrow W^+W^-$ in the same manner as the TGCs. The expected 95% confidence level limits for F_T for $\sqrt{s} = 500$ GeV and a luminosity of 500 fb^{-1} are shown in Figure 1, along with the predicted values of F_T for various masses M_ρ of a vector resonance in $W_L W_L$ scattering. The signal significances obtained by combining the results for $e^+e^- \rightarrow \nu\bar{\nu}W^+W^-$, $\nu\bar{\nu}ZZ$ [6, 7] with the F_T analysis of $e^+e^- \rightarrow W^+W^-$ [17] are displayed in Fig. 2 along with the results expected from the LHC [18]. At all values of the center-of-mass energy a linear collider provides a larger direct strong symmetry breaking signal than the LHC for vector resonance masses of 1200, 1600 and 2500 GeV. Only when the vector resonance disappears altogether (the LET case in the lower right-hand plot in Fig. 2) does the direct strong symmetry breaking signal from the $\sqrt{s} = 500$ GeV linear collider drop below the LHC signal. At higher e^+e^- center-of-mass energies the linear collider signal exceeds the LHC signal.




FIG. 1: 95% C.L. contour for F_T for $\sqrt{s} = 500$ GeV and 500 fb^{-1} . Values of F_T for various masses M_ρ of a vector resonance in $W_L W_L$ scattering are also shown. The F_T point “LET” refers to the case where no vector resonance exists at any mass in strong $W_L W_L$ scattering.

IV. STRONG WW SCATTERING BENCHMARK PROCESSES

The Snowmass 2001 working group on experimental approaches at linear colliders used a series of benchmarks to help evaluate the physics program of a future e^+e^- linear collider [19]. Strong WW scattering in the presence of scalar and vector resonances was simulated using the model of Han et al. [20], with resonance masses of




FIG. 2: Direct strong symmetry breaking signal significance in σ 's for various masses M_ρ of a vector resonance in $W_L W_L$ scattering. The numbers below the “LC” labels refer to the center-of-mass energy of the linear collider in GeV. The luminosity of the LHC is assumed to be 300 fb^{-1} , while the luminosities of the linear colliders are assumed to be 500, 1000, and 1000 fb^{-1} for $\sqrt{s}=500, 1000$, and 1500 GeV respectively. The lower right hand plot “LET” refers to the case where no vector resonance exists at any mass in strong $W_L W_L$ scattering.

1.0 and 1.5 TeV. The scalar resonance in this model was basically the SM Higgs. The widths of the vector resonances were 0.055 and 0.077 TeV for resonance masses of 1.0 and 1.5 TeV, respectively. For non-resonant strong WW scattering the unitarized K-matrix LET model[21] was used.

When estimating the mass scale reach of the K-matrix LET model and the mass resolution of the resonance model in the presence of a scalar ($I=0$) or tensor ($I=2$) resonance, we use the leading order modifications to the LET cross sections [22] :

$$\sigma(M_0) = \left(1 + \frac{8}{3} \frac{\hat{s}}{M_0^2}\right) \sigma_{\text{LET}}$$

Collider	Final State	\sqrt{s} TeV	\mathcal{L} fb ⁻¹	$M_0 = 1$ TeV	$M_0 = 1.5$ TeV	$M_1 = 1$ TeV		$M_1 = 1.5$ TeV	
				ΔM_0 GeV	ΔM_0 GeV	ΔM_1 GeV	$\Delta \Gamma_1$ GeV	ΔM_1 GeV	$\Delta \Gamma_1$ GeV
LC	W^+W^-	0.5	500	–	–	5.8	19.0	27.6	90
LC	W^+W^-	1.0	1000	89	249	0.01	0.03	4.0	13.5
LC	W^+W^-	1.5	1000	14	46	–	–	0.04	0.15

TABLE II: Expected error ΔM_0 for the mass of a scalar resonance, and expected errors ΔM_1 and $\Delta \Gamma_1$ for the mass and width, respectively, of a vector resonance. Results are shown for vector resonances of mass 1.0 and 1.5 TeV.

$$\sigma(M_2) = \left(1 + 2\frac{\hat{s}}{M_2^2}\right) \sigma_{\text{LET}} \quad ,$$

where M_0 and M_2 are the resonance masses in the $I = 0, 2$ channels, respectively. (The tensor resonance formula is used to estimate LHC mass scale sensitivity.) For detecting vector resonances we use the technipion form factor, which to leading order in s/M_1^2 is given by

$$F_T = \frac{M_1^2 - i\Gamma_1 M_1}{M_1^2 - s - i\Gamma_1 M_1} \quad ,$$

where M_1 and Γ_1 are the vector resonance mass and width, respectively. In order to evaluate the vector mass scale reach in the K-Matrix LET model we use the expression

$$Re(F_T) = 1 + \Delta_{\text{LET}} + \frac{s}{M_1^2} \quad ,$$

where Δ_{LET} is the contribution to F_T from strong WW scattering in the absence of a vector resonance. The dependence of Δ_{LET} on the details of the unitarization scheme grows as \sqrt{s} grows; the systematic uncertainty due to our lack of knowledge of these details is included in our calculations.

The expected errors for the mass of the scalar resonances are shown in Table II, along with the expected errors for the mass and width of the vector resonances. The measurement of the scalar mass M_0 is assumed to come solely from the measurement of the cross section σ , with $\sigma(M_0)$ defined above. For the measurement of the scalar mass there is a clear advantage in going to the higher CMS energy of 1.5 TeV. In contrast, the masses and widths of the vector resonances are measured very well at all CMS energies. Even the most poorly measured vector resonance parameter – the width of the 1.5 TeV resonance at $\sqrt{s} = 0.5$ TeV – is measured with an accuracy of 6% . At $\sqrt{s} = 1.0$ and 1.5 TeV the vector mass and width resolutions are typical of an e^+e^- collider sitting on top of the resonance.

Results for the K-matrix LET model are shown in Table III. The signal significance is displayed along with the 95% C.L. mass scale limits in the $I = 0, 1$ isospin channels. For comparison, results are also shown for the LHC in the $I = 2$ channel [18]. The tensor mass scale lower limit from the LHC is comparable to the scalar mass scale limits from the LC. Not suprisingly, the largest mass scale limits are the vector limits obtained in $e^+e^- \rightarrow W^+W^-$. Note that the vector mass scale lower limit M_1 does not improve as the CMS energy is raised from 1.0 to 1.5 TeV: this is due to the systematic uncertainty in the calculation of Δ_{LET} , which becomes important near $\sqrt{s} = 1.5$ TeV. The only way to reduce this particular systematic uncertainty is to actually do strong scattering experiments at the LHC and at an e^+e^- LC.

V. SUMMARY

Studies of strong electroweak symmetry breaking are enhanced by an e^+e^- linear collider with $\sqrt{s} = 0.5 - 1.5$ TeV. An LC complements a hadron collider nicely in providing better measurements of the chiral Lagrangian parameters L_{9L} and L_{9R} which affect triple gauge boson vertices. Also, the LC provides competitive measurements of the chiral Lagrangian parameters α_4 and α_5 which affect quartic gauge boson vertices.

A non-resonant strong symmetry breaking signal will be slightly larger at a $\sqrt{s} = 1.0$ TeV LC than at the LHC, and will be significantly larger if the e^+e^- CMS energy is raised to $\sqrt{s} = 1.5$ TeV. Less energy is required for strong vector resonance detection. A $\sqrt{s} = 0.5$ TeV LC provides a larger vector resonance signal than the

Collider	Final State	\sqrt{s} TeV	\mathcal{L} fb $^{-1}$	Signal signif.	M_0 (TeV) 95% C.L.	M_1 (TeV) 95% C.L.	M_2 (TeV) 95% C.L.
LC	W^+W^-	0.5	500	3σ	–	4.8	–
LC	W^+W^-	1.0	1000	7σ	–	6.4	–
LC	W^+W^-	1.5	1000	8σ	–	6.4	–
LC	$\nu\bar{\nu}W^+W^-, ZZ$	1.0	1000	7σ	1.7	–	–
LC	$\nu\bar{\nu}W^+W^-, ZZ$	1.5	1000	20σ	4.3	–	–
LHC	qqW^+W^+	14	300	9σ	–	–	3.0

TABLE III: Signal significance and 95% C.L. mass scale lower limits for the LET model with the K-matrix unitarization scheme. The mass scales M_0, M_1, M_2 correspond to structure in WW scattering in the $I=0,1$, and 2 isospin channels, respectively.

LHC for masses up to at least 2.5 TeV. The mass and width of a strong vector resonance can be measured at a LC with at least a few percent accuracy, even when the resonance lies well above the e^+e^- CMS energy.

Another important aspect of strong symmetry breaking is the study of $W^+W^- \rightarrow t\bar{t}$. This reaction can probably only be studied at a LC. Good strong symmetry breaking signals can be obtained in this channel at a LC, and these results should prove valuable in understanding electroweak symmetry breaking in the fermion sector.

Finally, we note that the systematic errors in signal and background calculations will be smaller at a LC than at a hadron collider, since the production mechanisms and backgrounds are limited to electroweak processes. However, we cannot at this time quantify this advantage since detailed studies of theoretical systematic errors in strong WW scattering have not been performed for either the LHC or the LC. This issue could be important given the size of some of the strong symmetry breaking signals and the paucity of sharp resonances in many strong symmetry breaking scenarios.

-
- [1] For recent reviews, see K. Lane, “Technicolor 2000,” hep-ph/0007304;
R. S. Chivukula and J. Womersley, in “Review of particle physics,” D. E. Groom *et al.*, Eur. Phys. J. **C15**, 1 (2000).
 - [2] R. Casalbuoni, A. Deandrea, S. De Curtis, D. Dominici, R. Gatto and J. F. Gunion, Nucl. Phys. B **555**, 3 (1999) [hep-ph/9809523].
 - [3] J. Bagger, S. Dawson, and G. Valencia, Nucl. Phys. **B399**, (1993) 364.
 - [4] T. Abe *et al.* [American Linear Collider Working Group Collaboration], SLAC-R-570 *Resource book for Snowmass 2001, 30 Jun - 21 Jul 2001, Snowmass, Colorado*.
 - [5] J. A. Bagger, A. F. Falk and M. Swartz Phys. Rev. Lett. **84**, 1385 (2000) [hep-ph/9908327].
 - [6] V. Barger, K. Cheung, T. Han and R. J. Phillips, Phys. Rev. D **52**, 3815 (1995) [hep-ph/9501379].
 - [7] E. Boos, H. J. He, W. Kilian, A. Pukhov, C. P. Yuan and P. M. Zerwas, Phys. Rev. D **57**, 1553 (1998) [hep-ph/9708310]; Phys. Rev. D **61**, 077901 (2000) [hep-ph/9908409].
 - [8] R. Chierici, S. Rosati, M. Kobel, “Strong electroweak symmetry breaking signals in WW scattering at TESLA.” LC-PHSM-2001-038. <http://www.desy.de/~lcnotes>
 - [9] T. L. Barklow, “Using $e^+e^- \rightarrow \nu\bar{\nu}t\bar{t}$ to Probe Strong Electroweak Symmetry Breaking at the NLC” in New Directions for High-Energy Physics: Snowmass 96, D.G. Cassel, L.T. Gennari, and R.H. Siemann, eds. (SLAC,1997).
 - [10] E. Ruiz Morales and M. E. Peskin, hep-ph/9909383.
 - [11] T. Han, Y. J. Kim, A. Likhoded and G. Valencia, Nucl. Phys. B **593**, 415 (2001) [hep-ph/0005306].
 - [12] F. Larios, T. M. Tait and C. P. Yuan, hep-ph/0101253.
 - [13] K. Hagiwara, R. D. Peccei, D. Zeppenfeld and K. Hikasa, Nucl. Phys. **B282** (1987) 253.
 - [14] A. Manohar and H. Georgi, Nucl. Phys. B **234**, 189 (1984).
 - [15] M. Peskin, in *Physics in Collisions IV*, A. Seiden, ed. (Éditions Frontières, Gif-Sur-Yvette, France, 1984).
 - [16] F. Iddir, A. Le Yaouanc, L. Oliver, O. Pene and J. C. Raynal, Phys. Rev. D **41**, 22 (1990).
 - [17] T. L. Barklow, “Strong symmetry breaking signals in $e^+e^- \rightarrow W^+W^-$ at $s^{**}(1/2) = 800\text{-GeV}$,” in Proceedings of 5th International Linear Collider Workshop (LCWS2000),Batavia, IL, October 24-28, 2000.
 - [18] ATLAS Detector and Physics Performance Technical Design Report, CERN LHCC 99-14/15 (1999).
 - [19] M. Battaglia, I. Hinchliffe, J. Jaros and J. Wells, “Experimental Approaches at Linear Colliders”, in these proceed-

ings.

- [20] T. Han, Y. J. Kim, A. Likhoded and G. Valencia, Nucl. Phys. B **593**, 415 (2001) [hep-ph/0005306].
- [21] M. S. Chanowitz and W. Kilgore, Phys. Lett. B **322**, 147 (1994) [hep-ph/9311336].
- [22] T. L. Barklow *et al.*, “Strong Coupling Electroweak Symmetry Breaking” in New Directions for High-Energy Physics: Snowmass 96, D.G. Cassel, L.T. Gennari, and R.H. Siemann, eds. (SLAC,1997). hep-ph/9704217.