

## **A Simplified Model of Intrabeam Scattering \***

K.L.F. Bane

*Stanford Linear Accelerator Center, Stanford University,  
Stanford, CA 94309 USA*

### *Abstract*

Beginning with the general Bjorken-Mtingwa solution, we derive a simplified model of intrabeam scattering (IBS), one valid for high energy beams in normal storage rings; our result is similar, though more accurate than a model due to Raubenheimer. In addition, we show that a modified version of Piwinski's IBS formulation (where  $\eta_{x,y}^2/\beta_{x,y}$  has been replaced by  $\mathcal{H}_{x,y}$ ) at high energies asymptotically approaches the same result.

*Presented at the Eighth European Particle Accelerator Conference (EPAC'02),  
Paris, France  
June 3-7, 2002*

---

\*Work supported by Department of Energy contract DE-AC03-76SF00515.

# A SIMPLIFIED MODEL OF INTRABEAM SCATTERING

K.L.F. Bane, SLAC, Stanford, CA94309, USA \*

## Abstract

Beginning with the general Bjorken-Mtingwa solution, we derive a simplified model of intrabeam scattering (IBS), one valid for high energy beams in normal storage rings; our result is similar, though more accurate than a model due to Raubenheimer. In addition, we show that a modified version of Piwinski's IBS formulation (where  $\eta_{x,y}^2/\beta_{x,y}$  has been replaced by  $\mathcal{H}_{x,y}$ ) at high energies asymptotically approaches the same result.

## 1 INTRODUCTION

Intrabeam scattering (IBS), an effect that tends to increase the beam emittance, is important in hadronic[1] and heavy ion[2] circular machines, as well as in low emittance electron storage rings[3]. In the former type of machines it results in emittances that continually increase with time; in the latter type, in steady-state emittances that are larger than those given by quantum excitation/synchrotron radiation alone.

The theory of intrabeam scattering for accelerators was first developed by Piwinski[4], a result that was extended by Martini[5], to give a formulation that we call here the standard Piwinski (P) method[6]; this was followed by the equally detailed Bjorken and Mtingwa (B-M) result[7]. Both approaches solve the local, two-particle Coulomb scattering problem for (six-dimensional) Gaussian, uncoupled beams, but the two results appear to be different; of the two, the B-M result is thought to be the more general[8].

For both the P and the B-M methods solving for the IBS growth rates is time consuming, involving, at each time (or iteration) step, a numerical integration at every lattice element. Therefore, simpler, more approximate formulations of IBS have been developed over the years: there are approximate solutions of Parzen[9], Le Duff[10], Raubenheimer[11], and Wei[12]. In the present report we derive—starting with the general B-M formalism—another approximation, one valid for high energy beams and more accurate than Raubenheimer's approximation. We, in addition, demonstrate that under these same conditions a modified version of Piwinski's IBS formulation asymptotically becomes equal to this result.

## 2 HIGH ENERGY APPROXIMATION

### 2.1 The General B-M Solution[7]

Let us consider bunched beams that are uncoupled, and include vertical dispersion due to *e.g.* orbit errors. Let the

intrabeam scattering growth rates be

$$\frac{1}{T_p} = \frac{1}{\sigma_p} \frac{d\sigma_p}{dt}, \quad \frac{1}{T_x} = \frac{1}{\epsilon_x^{1/2}} \frac{d\epsilon_x^{1/2}}{dt}, \quad \frac{1}{T_y} = \frac{1}{\epsilon_y^{1/2}} \frac{d\epsilon_y^{1/2}}{dt}, \quad (1)$$

with  $\sigma_p$  the relative energy spread,  $\epsilon_x$  the horizontal emittance, and  $\epsilon_y$  the vertical emittance. The growth rates according to Bjorken-Mtingwa (including a  $\sqrt{2}$  correction factor[13], and including vertical dispersion) are

$$\frac{1}{T_i} = 4\pi A(\log) \left\langle \int_0^\infty \frac{d\lambda \lambda^{1/2}}{[\det(L + \lambda I)]^{1/2}} \left\{ Tr L^{(i)} Tr \left( \frac{1}{L + \lambda I} \right) - 3 Tr L^{(i)} \left( \frac{1}{L + \lambda I} \right) \right\} \right\rangle \quad (2)$$

where  $i$  represents  $p$ ,  $x$ , or  $y$ ;

$$A = \frac{r_0^2 c N}{64\pi^2 \beta^3 \gamma^4 \epsilon_x \epsilon_y \sigma_s \sigma_p}, \quad (3)$$

with  $r_0$  the classical particle radius,  $c$  the speed of light,  $N$  the bunch population,  $\beta$  the velocity over  $c$ ,  $\gamma$  the Lorentz energy factor, and  $\sigma_s$  the bunch length;  $(\log)$  represents the Coulomb log factor,  $\langle \rangle$  means that the enclosed quantities, combinations of beam parameters and lattice properties, are averaged around the entire ring;  $\det$  and  $Tr$  signify, respectively, the determinant and the trace of a matrix, and  $I$  is the unit matrix. Auxiliary matrices are defined as

$$L = L^{(p)} + L^{(x)} + L^{(y)}, \quad (4)$$

$$L^{(p)} = \frac{\gamma^2}{\sigma_p^2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (5)$$

$$L^{(x)} = \frac{\beta_x}{\epsilon_x} \begin{pmatrix} 1 & -\gamma\phi_x & 0 \\ -\gamma\phi_x & \gamma^2 \mathcal{H}_x / \beta_x & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (6)$$

$$L^{(y)} = \frac{\beta_y}{\epsilon_y} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \gamma^2 \mathcal{H}_y / \beta_y & -\gamma\phi_y \\ 0 & -\gamma\phi_y & 1 \end{pmatrix}. \quad (7)$$

The dispersion invariant is  $\mathcal{H} = [\eta^2 + (\beta\eta' - \frac{1}{2}\beta'\eta)^2]/\beta$ , and  $\phi = \eta' - \frac{1}{2}\beta'\eta/\beta$ , where  $\beta$  and  $\eta$  are the beta and dispersion lattice functions.

### The Bjorken-Mtingwa Solution at High Energies

Let us first consider  $1/T_p$  as given by Eq. 2. Note that if we change the integration variable to  $\lambda' = \lambda\sigma_H^2/\gamma^2$  then

$$(L + \lambda' I) = \frac{\gamma^2}{\sigma_H^2} \begin{pmatrix} a^2 + \lambda' & -a\zeta_x & 0 \\ -a\zeta_x & 1 + \lambda' & -b\zeta_y \\ 0 & -b\zeta_y & b^2 + \lambda' \end{pmatrix}, \quad (8)$$

\* Work supported by the Department of Energy, contract DE-AC03-76SF00515

with

$$\frac{1}{\sigma_H^2} = \frac{1}{\sigma_p^2} + \frac{\mathcal{H}_x}{\epsilon_x} + \frac{\mathcal{H}_y}{\epsilon_y} \quad , \quad (9)$$

$$a = \frac{\sigma_H}{\gamma} \sqrt{\frac{\beta_x}{\epsilon_x}} \quad , \quad b = \frac{\sigma_H}{\gamma} \sqrt{\frac{\beta_y}{\epsilon_y}} \quad , \quad \zeta_{x,y} = \phi_{x,y} \sigma_H \sqrt{\frac{\beta_{x,y}}{\epsilon_{x,y}}} \quad (10)$$

Note that, other than a multiplicative factor, there are only 4 parameters in this matrix:  $a, b, \zeta_x, \zeta_y$ . Note that, since  $\beta\phi^2 \leq \mathcal{H}$ , the parameters  $\zeta < 1$ ; and that if  $\mathcal{H} \approx \eta^2/\beta$  then  $\zeta$  is small. We give, in Table 1, average values of  $a, b, \zeta_x$ , in selected electron rings.

Table 1: Average values of  $a, b, \zeta_x$ , in selected electron rings. The zero current emittance ratio  $\sim 0.5\%$  in all cases.

| Machine   | $E[\text{GeV}]$ | $N[10^{10}]$ | $\langle a \rangle$ | $\langle b \rangle$ | $\langle \zeta_x \rangle$ |
|-----------|-----------------|--------------|---------------------|---------------------|---------------------------|
| KEK's ATF | 1.4             | .9           | .01                 | .10                 | .15                       |
| NLC       | 2.0             | .75          | .01                 | .20                 | .40                       |
| ALS       | 1.0             | 5.           | .015                | .25                 | .15                       |

Let us limit consideration to high energies, specifically let us assume  $a, b \ll 1$  (if the beam is cooler longitudinally than transversely, then this is satisfied). We note that all 3 rings in Table 1, on average, satisfy this condition reasonably well. Assuming this condition, the 2nd term in the braces of Eq. 2 is small compared to the first term, and we drop it. Our second assumption is to drop off-diagonal terms (let  $\zeta = 0$ ), and then all matrices will be diagonal.

Simplifying the remaining integral by applying the high energy assumption we finally obtain

$$\frac{1}{T_p} \approx \frac{r_0^2 c N (\log)}{16 \gamma^3 \epsilon_x^{3/4} \epsilon_y^{3/4} \sigma_s \sigma_p^3} \left\langle \sigma_H g(a/b) (\beta_x \beta_y)^{-1/4} \right\rangle \quad , \quad (11)$$

with

$$g(\alpha) = \frac{2\sqrt{\alpha}}{\pi} \int_0^\infty \frac{du}{\sqrt{1+u^2} \sqrt{\alpha^2+u^2}} \quad . \quad (12)$$

A plot of  $g(\alpha)$  over the interval  $[0 < \alpha < 1]$  is given in Fig. 1; to obtain the results for  $\alpha > 1$ , note that  $g(\alpha) = g(1/\alpha)$ . A fit to  $g$ ,

$$g(\alpha) \approx \alpha^{(0.021-0.044 \ln \alpha)} \quad [\text{for } 0.01 < \alpha < 1] \quad , \quad (13)$$

is given by the dashes in Fig. 1. The fit has a maximum error of 1.5% over  $[0.02 \leq \alpha \leq 1]$ .

Similarly, beginning with the 2nd and 3rd of Eqs. 2, we obtain

$$\frac{1}{T_{x,y}} \approx \frac{\sigma_p^2 \langle \mathcal{H}_{x,y} \rangle}{\epsilon_{x,y}} \frac{1}{T_p} \quad . \quad (14)$$

Our approximate IBS solution is Eqs. 11,14. Note that Parzen's high energy formula is a similar, though more approximate, result to that given here[9]; and Raubenheimer's approximation is Eq. 11, with  $g(a/b)\sigma_H/\sigma_p$  replaced by  $\frac{1}{2}$ , and Eqs. 14 exactly as given here[11].

Note that the beam properties in Eqs. 11,14, need to be the self-consistent values. Thus, for example, to find the




Figure 1: The auxiliary function  $g(\alpha)$  (solid curve) and the approximation,  $g = \alpha^{(0.021-0.044 \ln \alpha)}$  (dashes).

steady-state growth rates in electron machines, iteration will be required[6]. Note also that these equations assume that the zero-current vertical emittance is due mainly to vertical dispersion caused by orbit errors; if it is due mainly to (weak)  $x$ - $y$  coupling we let  $\mathcal{H}_y = 0$ , drop the  $1/T_y$  equation, and let  $\epsilon_y = \kappa \epsilon_x$ , with  $\kappa$  the coupling factor[3].

What sort of error does our model produce? Consider a position in the ring where  $\zeta_y = 0$ . In Fig. 2 we plot the ratio of the *local* growth rate  $T_p^{-1}$  as given by our model to that given by Eq. 2 as function of  $\zeta_x$ , for example combinations of  $a$  and  $b$ . We see that for  $\zeta_x \lesssim \sqrt{b}e^{(1-\sqrt{b})}$  (which is typically true in storage rings) the dependance on  $\zeta_x$  is weak and can be ignored. In this region we see that the model approaches B-M from above as  $a, b \rightarrow 0$ . Finally, adding small  $\zeta_y \neq 0$  will reduce slightly the ratio of Fig. 2.




Figure 2: The ratio of *local* growth rates in  $p$  as function of  $\zeta_x$ , for  $b = 0.1$  (blue) and  $b = 0.2$  (red) [ $\zeta_y = 0$ ].

### 3 COMPARISON TO PIWINSKI

#### 3.1 The Standard Piwinski Solution[6]

The standard Piwinski solution is

$$\frac{1}{T_p} = A \left\langle \frac{\sigma_h^2}{\sigma_p^2} f(\tilde{a}, \tilde{b}, q) \right\rangle$$

$$\frac{1}{T_x} = A \left\langle f\left(\frac{1}{\tilde{a}}, \frac{\tilde{b}}{\tilde{a}}, \frac{q}{\tilde{a}}\right) + \frac{\eta_x^2 \sigma_h^2}{\beta_x \epsilon_x} f(\tilde{a}, \tilde{b}, q) \right\rangle$$

$$\frac{1}{T_y} = A \left\langle f\left(\frac{1}{\tilde{b}}, \frac{\tilde{a}}{\tilde{b}}, \frac{q}{\tilde{b}}\right) + \frac{\eta_y^2 \sigma_h^2}{\beta_y \epsilon_y} f(\tilde{a}, \tilde{b}, q) \right\rangle ; \quad (15)$$

$$\frac{1}{\sigma_h^2} = \frac{1}{\sigma_p^2} + \frac{\eta_x^2}{\beta_x \epsilon_x} + \frac{\eta_y^2}{\beta_y \epsilon_y} , \quad (16)$$

$$\tilde{a} = \frac{\sigma_h}{\gamma} \sqrt{\frac{\beta_x}{\epsilon_x}}, \quad \tilde{b} = \frac{\sigma_h}{\gamma} \sqrt{\frac{\beta_y}{\epsilon_y}}, \quad q = \sigma_h \beta \sqrt{\frac{2d}{r_0}} ; \quad (17)$$

the function  $f$  is given by:

$$f(\tilde{a}, \tilde{b}, q) = 8\pi \int_0^1 du \frac{1 - 3u^2}{PQ} \times \left\{ 2 \ln \left[ \frac{q}{2} \left( \frac{1}{P} + \frac{1}{Q} \right) \right] - 0.577 \dots \right\} \quad (18)$$

$$P^2 = \tilde{a}^2 + (1 - \tilde{a}^2)u^2, \quad Q^2 = \tilde{b}^2 + (1 - \tilde{b}^2)u^2 . \quad (19)$$

The parameter  $d$  functions as a maximum impact parameter, and is normally taken as the vertical beam size.

### 3.2 Comparison of Modified Piwinski to the B-M Solution at High Energies

We note that Piwinski's result depends on  $\eta^2/\beta$ , and not on  $\mathcal{H}$  and  $\phi$ , as the B-M result does. This may suffice for rings with  $\langle \mathcal{H} \rangle \approx \langle \eta^2/\beta \rangle$ . For a general comparison, however, let us consider a formulation that we call the *modified* Piwinski solution. It is the standard version of Piwinski, but with  $\eta^2/\beta$  replaced by  $\mathcal{H}$  (*i.e.*  $\tilde{a}$ ,  $\tilde{b}$ ,  $\sigma_h$ , become  $a$ ,  $b$ ,  $\sigma_H$ , respectively).

Let us consider high energy beams, *i.e.* let  $a, b \ll 1$ : First, notice that in the integral of the auxiliary function  $f$  (Eq. 18): the  $-0.577$  can be replaced by 0; the  $-3u^2$  in the numerator can be set to 0;  $P$  ( $Q$ ) can be replaced by  $\sqrt{a^2 + u^2}$  ( $\sqrt{b^2 + u^2}$ ). The first term in the braces can be approximated by a constant and then be pulled out of the integral; it becomes the effective Coulomb log factor. Note that for the proper choice of the Piwinski parameter  $d$ , the effective Coulomb log can be made the same as the B-M parameter (log). For flat beams ( $a \ll b$ ), the Coulomb log of Piwinski becomes  $(\log) = \ln[d\sigma_H^2/(4r_0a^2)]$ .

We finally obtain, for the first of Eqs. 15,

$$\frac{1}{T_p} \approx \frac{r_0^2 c N (\log)}{16 \gamma^3 \epsilon_x^{3/4} \epsilon_y^{3/4} \sigma_s \sigma_p^3} \left\langle \sigma_H h(a, b) (\beta_x \beta_y)^{-1/4} \right\rangle , \quad (20)$$

with

$$h(a, b) = \frac{2\sqrt{ab}}{\pi} \int_0^1 \frac{du}{\sqrt{a^2 + u^2} \sqrt{b^2 + u^2}} . \quad (21)$$

We see that the approximate equation for  $1/T_p$  for high energy beams according to modified Piwinski is the same as that for B-M, except that  $h(a, b)$  replaces  $g(a/b)$ . But for  $a, b$  small,  $h(a, b) \approx g(a/b)$ , and the Piwinski result approaches the B-M result. For example, for the ATF with  $\epsilon_y/\epsilon_x \sim 0.01$ ,  $a \sim 0.01$ ,  $a/b \sim 0.1$ , and  $h(a, b)/g(a/b) = 0.97$ ; the agreement is quite good.

Finally, for the relation between the transverse to longitudinal growth rates according to modified Piwinski: note that for non-zero vertical dispersion the second term in the brackets of Eqs. 15 (but with  $\eta_{x,y}^2/\beta_{x,y}$  replaced by  $\mathcal{H}_{x,y}$ ), will tend to dominate over the first term, and the results become the same as for the B-M method.

In summary, we have shown that for high energy beams ( $a, b \ll 1$ ), in normal rings ( $\zeta$  not very close to 1): if the parameter  $d$  in P is chosen to give the same equivalent Coulomb log as in B-M, then the *modified* Piwinski solution agrees with the Bjorken-Mtingwa solution.

## 4 NUMERICAL COMPARISON[3]

We consider a numerical comparison between results of the general B-M method, the modified Piwinski method, and Eqs. 11,14. The example is the ATF ring with no coupling; to generate vertical errors, magnets were randomly offset by  $15 \mu\text{m}$ , and the closed orbit was found. For this example  $\langle \mathcal{H}_y \rangle = 17 \mu\text{m}$ , yielding a zero-current emittance ratio of 0.7%; the beam current is 3.1 mA. The steady-state growth rates according to the 3 methods are given in Table 2. We note that the Piwinski results are 4.5% low, and the results of Eqs. 11,14, agree very well with those of B-M. Additionally, note that, not only the (averaged) growth rates, but even the *local* growth rates around the ring agree well for the three cases. Finally, note that for coupling dominated NLC, ALS examples ( $\kappa = 0.5\%$ , see Table 1) the error in the steady-state growth rates ( $T_p^{-1}, T_x^{-1}$ ) obtained with the model is (12%, 2%), (7%, 0%), respectively.

Table 2: Steady-state IBS growth rates (in  $\text{s}^{-1}$ ) for an ATF example with vertical dispersion due to random errors.

| Method            | $1/T_p$ | $1/T_x$ | $1/T_y$ |
|-------------------|---------|---------|---------|
| Modified Piwinski | 25.9    | 24.7    | 18.5    |
| Bjorken-Mtingwa   | 27.0    | 26.0    | 19.4    |
| Eqs. 11,14        | 27.4    | 26.0    | 19.4    |

The author thanks A. Piwinski, K. Kubo and other coauthors of Ref. [3] for help in understanding IBS theory; K. Kubo, A. Wolski, C. Steier, for supplying the lattices of the ATF, NLC, ALS rings, respectively.

## 5 REFERENCES

- [1] C. Bhat, *et al*, *Proc. PAC99*, New York (1999) 3155.
- [2] W. Fischer, *et al*, *Proc. PAC2001*, Chicago (2001) 2857.
- [3] K. Bane, *et al*, SLAC-PUB-9227, May 2002.
- [4] A. Piwinski, Tech. Rep. HEAC 74, Stanford, 1974.
- [5] M. Martini, Tech. Rep. PS/84-9(AA), CERN, 1984.
- [6] A. Piwinski, in *Handbook of Accelerator Physics*, World Scientific (1999) 125.
- [7] J. Bjorken and S. Mtingwa, *Part. Accel.*, **13** (1983) 115.
- [8] A. Piwinski, private communication.
- [9] G. Parzen, *Nucl. Instr. Meth.*, **A256** (1987) 231.
- [10] J. Le Duff, *Proc. of CERN Accel. School* (1989) 114.
- [11] T. Raubenheimer, SLAC-R-387, PhD thesis, 1991, Sec. 2.3.1.
- [12] J. Wei, *Proc. PAC93*, Washington, D.C. (1993) 3651.
- [13] K. Kubo and K. Oide, *PRST-AB*, **4** (2001) 124401.