

## Experimental characterization of nonlinear harmonic radiation from a visible SASE FEL at saturation

A. Tremaine<sup>1</sup>, X.J. Wang<sup>2</sup>, M. Babzien<sup>2</sup>, I. Ben-Zvi<sup>2</sup>, M. Cornacchia<sup>3</sup>,  
A. Murokh<sup>1</sup>, H.-D. Nuhn<sup>3</sup>, R. Malone<sup>2</sup>, C. Pellegrini<sup>1</sup>, S. Reiche<sup>1</sup>,  
J. Rosenzweig<sup>1</sup>, J. Skaritka<sup>2</sup> and V. Yakimenko<sup>2</sup>

<sup>1</sup>*Department of Physics & Astronomy, UCLA, Los Angeles, CA 90095*

<sup>2</sup>*Accelerator Test Facility, NSLS, BNL, Upton, NY 11973*

<sup>3</sup>*SSRL, SLAC, Stanford, CA 94309*

Nonlinear harmonic radiation was observed using the VISA SASE FEL at saturation. The gain lengths, spectra and energies of the three lowest SASE FEL modes were experimentally characterized. Both the measured harmonic gain lengths and center spectral wavelengths are shown to decrease with harmonic number,  $n$ , which is consistent with nonlinear harmonic theory. The measured energies for both second and third harmonics are about 1% of the fundamental; the strong second harmonic radiation ( $\approx 1 \mu\text{J}$ ) observed is unique for low energy SASE FELs. These experimental results demonstrate for the first time the feasibility of using nonlinear harmonic SASE FEL radiation to produce coherent, femtosecond X-rays.

*PACS* numbers: 41.60.Cr, 41.60.Ap, 41.85.Ja

Submitted to *Physical Review Letters*

The explosive growth of synchrotron light source applications in the last few decades is due largely to the improved brightness of the X-rays beams they produce [1]. During the same time, high power, femtosecond laser technologies have opened new frontiers in spectroscopy and dynamics system studies [2]. The free electron laser (FEL) is one of the technologies that will bridge the gap between laser light (coherent, femtosecond pulse duration at long wavelengths) and synchrotron light (incoherent, picosecond pulse duration at X-ray wavelengths.)

An FEL can operate either as an oscillator or a single-pass amplifier. For coherent, femtosecond X-ray generation, oscillators and laser-seeded FEL amplifiers are not feasible due to the non-existence of highly reflective mirrors and laser seeds in the X-ray region. All proposed X-ray FEL facilities [3,4] are based on the SASE FEL, which is a high-gain, single-pass FEL amplifier that starts from noise [5,6]. SASE X-ray facilities rely on high-brightness, low energy-spread electron beams and novel undulators to achieve saturation within a reasonable distance.

One of the natural consequences of the SASE FEL is nonlinear harmonic generation (NHG) [7-10], which occurs only in the high gain and saturation regime. Compared to other harmonic radiation, NHG is driven by the fundamental mode, and has faster growth rates and higher intensities. Both simulation and theory predict the NHG 3<sup>rd</sup> harmonic to be about 1% of the fundamental for an X-ray SASE FEL [9-10]. This is about eight orders of magnitude brighter than current third generation synchrotron light sources. NHG is transversely coherent, and it will make it possible for SASE FELs to have broad spectral coverage at minimum cost. In addition, NHG is less sensitive to the electron beam parameters, undulator errors and wake fields compared to other linear harmonic generation schemes [11].

As an electron beam passes through the periodic magnetic field of an undulator, such as in a SASE FEL, spontaneous radiation is emitted due to the electron beam/magnetic field interaction. A ponderomotive potential is formed by the interaction of this radiation field with the undulator magnetic field. The ponderomotive potential longitudinally microbunches the electron beam on the scale of a radiation wavelength,  $\lambda_r$ . As the beam microbunches further, the coherent radiation emitted from the beam grows exponentially through the undulator. At saturation, the radiation field stops

growing and an oscillatory exchange of energy between the radiation field and electron beam continues until the end of the undulator.

The wavelength of fundamental radiation emitted from this process is given by

$$\lambda_r = \frac{\lambda_u}{2\gamma^2} \left( 1 + \frac{K^2}{2} \right) \quad (1)$$

where  $\lambda_u$  is the undulator period,  $\gamma$  is the electron beam energy, and  $K$  is the undulator parameter. Eq. 1 is also the period at which the electron beam is longitudinally microbunched,  $\lambda_{mb} = \lambda_r$ . High-gain SASE FELs have been demonstrated from IR to UV wavelengths [12-22], and saturation of SASE FELs has been observed recently at the LEUTL [19], VISA [20-21] and TTF [22] facilities. In addition, measurements of the longitudinal microbunching of the electron beam from SASE FELs have been reported [23-24].

An electron beam in an FEL near saturation is tightly microbunched at the fundamental wavelength. Such a strongly microbunched electron beam drives the microbunching at its harmonics [8],  $\lambda_{mb,n} = \lambda_{mb} / n$  where  $n = 1, 2, 3, \dots$ . Since the harmonic microbunching is driven by the fundamental ( $n=1$ ) SASE process, the harmonic microbunching grows at a rate faster than the fundamental, but starts later in the undulator. The radiation emitted from such a bunching process is known as nonlinear harmonic radiation [8-10]. The experimental investigation of the nonlinear harmonic radiation at the VISA SASE FEL is presented in this letter. For the first time, the gain lengths for the three lowest modes of a SASE FEL have been measured and shown to decrease with the harmonic number,  $n$ . The measured energies from nonlinear harmonic generation are shown to agree with both nonlinear theory and simulations. In addition, the spectra for these modes have been experimentally characterized.

The VISA FEL is part of the R&D effort for the proposed LCLS X-ray FEL project [3] and is a test bed for both the high-brightness electron beam and SASE FEL physics required for an X-ray FEL. Table I summarizes the key design parameters for the experiment.

Table I. VISA FEL design parameters

Wavelength (Fundamental)	$\lambda_f$	800 nm
E-beam Energy	E	72 MeV
Peak Current	$I_p$	200 A
Emittance	$\varepsilon$	2 mm-mrad
Energy spread	$\Delta\gamma/\gamma$	.17 %
Power Gain Length	$L_g$	17.5 cm
Spot size	$\sigma$	60 $\mu\text{m}$

The VISA experiment was carried out at the Accelerator Test Facility (ATF) of Brookhaven National Laboratory. The ATF high-brightness electron beam is produced by an S-band, 1.6-cell photocathode RF gun operating in longitudinal emittance compensation mode [26]. The electron beam is accelerated to 72 MeV by two sections of traveling wave linac. The final match of the electron beam into the VISA undulator is done by a quadrupole magnet triplet. The VISA undulator [27] is a planar magnet array with built-in strong focusing. The strong focusing is introduced by placing dipole magnets with opposite polarity on either side of the electron beam propagation axis [27]. The 4 m undulator has a 1.8 cm period and a 6 mm gap. The combination of strong focusing and bright electron beam demands the straightness of the undulator to be better than 30  $\mu\text{m}$  over one gain length [28].

Both SASE and NHG measurements along the undulator are made using eight diagnostic pop-in ports [29] plus the diagnostic station downstream of the undulator. The diagnostic pop-in screens inside the undulator are spaced at 50 cm intervals, the last pop-in being 25 cm from the exit of the undulator. Nonlinear harmonic growth occurs primarily in the last quarter of the undulator and can be measured at the last three diagnostic pop-ins. The fundamental SASE radiation can be detected at the last five diagnostics ports.

The nonlinear harmonic spectral measurements in Fig. 1(a) were captured in a single shot when high SASE gain was established. The spectrometer (Ocean Optics) has peak sensitivity around 300 nm and attenuates (by about two orders of magnitude) at the

SASE fundamental. The resolution of the spectrometer is about 1 nm. Fig. 1(a) shows the harmonic wavelengths, decreasing according to the following equation:

$$\lambda_{r,n} = \frac{\lambda_f}{n} \quad (2)$$

where  $n = 1, 2, 3, \dots$  is the harmonic number and  $\lambda_f$  is the fundamental wavelength ( $n = 1$ ). The measured fundamental wavelength is 845 nm while the 2<sup>nd</sup> and the 3<sup>rd</sup> harmonics are 423 nm and 281 nm ( $n = 2, 3$ ), respectively. The spectrum obtained in Fig. 1(a) agrees with Eq. 2 within the resolution of the spectrometer.

Fig. 1 also shows the spectral bandwidth,  $\Delta\lambda_n$  (FWHM), which decreases as the mode number increases. The bandwidth measured for the fundamental at saturation has been reported elsewhere for VISA [21-22], and is  $\Delta\lambda_f = 15$  nm. Fig. 1(b) provides independent measurements for the 2<sup>nd</sup> and 3<sup>rd</sup> harmonics at the VISA saturation. The spectral widths (FWHM) for the 2<sup>nd</sup> and 3<sup>rd</sup> harmonic are  $\Delta\lambda_2 = 6$  nm and  $\Delta\lambda_3 = 3$  nm, respectively. The above data shows the nonlinear harmonic bandwidths,  $\Delta\lambda_n$ , relative to that of the fundamental decrease with mode number,  $n$ . In addition, the relative bandwidths,  $\Delta\lambda_n / \lambda_n$ , also decrease with mode number, but at a slower rate. Recent studies [30] of the SASE nonlinear harmonic spectra support this observation. A double hump was observed in Fig. 1(a) for the 2<sup>nd</sup> harmonic. Fig. 1(a) was captured when the nonlinear harmonics were just beginning to grow and the double hump is most probably due to the electron beam/harmonic radiation interaction during the SASE NHG start-up process. Figure 1 experimentally confirms that high quality nonlinear harmonic spectra can be expected for future short wavelength SASE FELs driven by high-brightness electron beams.

The nonlinear harmonics as a function of distance for the three lowest SASE FEL modes were measured using factory-calibrated Molelectron joulemeters. Two bandpass filters were placed before the joulemeter for each harmonic measurement. Each filter attenuates the fundamental signal by at least a factor of 1000, thus combining two filters leads to six orders of magnitude attenuation outside of the pass-band. Fig. 2 shows a log-linear plot for fundamental, 2<sup>nd</sup> and 3<sup>rd</sup> harmonics vs. distance along the second half of

the 4-m VISA undulator. The deviation of the last two points from exponential growth for the fundamental mode demonstrates SASE saturation. The fundamental saturation length is about 3.8 m.

The fundamental gain length of  $L_g = 19$  cm is obtained from Fig. 2. To obtain the harmonic gain lengths,  $L_{g,2}$  and  $L_{g,3}$ , the data in the linear harmonic region are excluded. The 2<sup>nd</sup> harmonic gain length is measured to be  $L_{g,2} = 9.8$  cm. Because of the limited number of diagnostic pop-ins, only two points for the nonlinear 3<sup>rd</sup> harmonic regime could be used to estimate the gain length. Multiple measurements resulted in a 6 cm average growth rate, which is consistent with the gain length relation quoted in Eq. 4, below. Figure 2 shows the nonlinear harmonic radiation grows faster compared to the fundamental. Our measurements confirm nonlinear harmonic generation theory [8-10], in which the gain length decreases with the harmonic number,  $n$ .

$$L_{g,n} = \frac{L_g}{n}, \quad (4)$$

Finally, the energy for each mode was measured downstream of the undulator. Table II shows the nonlinear harmonic energies measured at the end of the

Table II. Measured Harmonic Energy for fundamental, 2<sup>nd</sup> and 3<sup>rd</sup> harmonics

Mode	n	1	2	3
Wavelength (nm)	$\lambda_f$	845	423	281
Energy ( $\mu$ J)	$E_n$	52	.93	.40
% of fundamental energy	$\frac{E_n}{E_1} \times 100$		1.8	0.77

undulator and the harmonic energy as a percentage of the fundamental. It can be seen from Table II that not only is there significant UV (281 nm, 3<sup>rd</sup> harmonic) energy, but also comparable energy in the green (423 nm, 2<sup>nd</sup> harmonic). The relatively large amount

of 2<sup>nd</sup> harmonic energy from a planar undulator is due to the low electron beam energy used in this experiment. The 2<sup>nd</sup> and 3<sup>rd</sup> nonlinear harmonic energies are related by [31],

$$E_2 \approx E_3 \left( \frac{K}{\gamma k_u \sigma_x} \right)^2 \left( \frac{K_2}{K_3} \right)^2 \left( \frac{b_2}{b_3} \right)^2 \quad (5)$$

where  $K=1.28$  is the undulator parameter,  $\gamma$  is the beam energy,  $k_u$  is the undulator wave number,  $\sigma_x$  is the RMS beam size, and  $b_n$  are the harmonic bunching parameters.  $K_n$  are the coupling coefficients [31]:

$$K_n = K(-1)^{(n-1)/2} [J_{(n-1)/2}(n\xi) - J_{(n+1)/2}(n\xi)], \quad n = 1, 3, 5 \dots \quad (6)$$

$$K_n = K(-1)^{(n-2)/2} [J'_{n/2}(n\xi)], \quad n = 2, 4, 6 \dots$$

where  $\xi = K^2 / (4 + 2K^2)$  and  $J(n\xi)$  is the Bessel function.

Using the parameters in Table I, Genesis [32] simulations produce harmonic bunching parameters of  $b_2 = .47$ , and  $b_3 = .28$  near saturation. Inserting the GENESIS results into Eqs. 5 and 6, NHG theory predicts that the 3<sup>rd</sup> and 2<sup>nd</sup> harmonic energies should be 1% and 1.6% of the fundamental, respectively. The measured 3<sup>rd</sup> and 2<sup>nd</sup> harmonics are 0.8 and 1.8% (Table 2) of the fundamental, with the ratio  $E_2 / E_3 = 2.3$ .

The slight discrepancy between the measured and calculated energy ratio could arise from two sources: First, for each harmonic measurement, the signal passes through two narrow band filters. The errors in calibrating the transmission coefficient for such filters is one possible source, especially for the 3<sup>rd</sup> harmonic where each filter attenuates the desired signal by 80%. Second, the VISA beam profile monitors have a resolution no better than 15  $\mu\text{m}$ . An error in spot size measurements of this order could account for the remainder of the discrepancy described above since the energy ratio given in Eq. 5 strongly depends on the ratio of the wiggles amplitude to the transverse spot size,  $K / \gamma k_u : \sigma_x$ .

The on-axis radiation is dominated by the odd harmonics for a planar undulator. If the off-axis radiation is collected, the even harmonic is observed to be quite significant

and comparable to the odd harmonic for the VISA FEL. Coupling to the 2<sup>nd</sup> harmonic strengthens as the electron beam energy decreases. The reason for this can be seen as follows: If the electron beam energy decreases, then the wiggler amplitude will increase. The even harmonics for a planar undulator are emitted off-axis [33], and as the transverse wiggler amplitude increases relative to  $\sigma_x$ , so will the radiation emitted for the 2<sup>nd</sup> harmonic. The strong coupling to even the NHG occurs only for low energy FELs, such as VISA, thus providing a unique window to further test the NHG theory.

Though the observation of NHG in the far infrared is claimed [34], we have presented the first experimental characterization of the nonlinear harmonic gain lengths and single shot spectra for the three lowest modes of a SASE FEL. The basic properties of NHG theory have been experimentally confirmed; gain length and harmonic wavelength decrease linearly with harmonic number,  $n$ , and harmonic spectral width narrows with harmonic number,  $n$ . Furthermore, large energies of the 2<sup>nd</sup> and 3<sup>rd</sup> nonlinear harmonics were observed and confirmed the predictions of NHG theory. These results are important not only as a comprehensive verification of nonlinear harmonic SASE FEL theory, but also as a demonstration of an alternative source of high-energy, high-quality spectra, short wavelength radiation.

The authors would like to thank Dr. Z. Huang of the APS for stimulating conversations regarding the nonlinear harmonic evolution process. We would also like to express our gratitude to Mr. R. Harrington and Mr. D. Davis for their support. This work is supported by US Department of Energy contracts DE-AC02-98CH10886, DE-FG03-92ER40793, and DE-AC03-76F00515.

#### References:

- [1] J.-L. Laclare, Light Source Performance Achievements, in *Proceeding of 1998 European Particle Accelerator Conference*, 79 (1998).



- [2] A. H. Zewail, *Femtochemistry: Dynamics with Atomic Resolution*, Femtochemistry & Femtobiology, V. Sundström (Ed.), World Scientific, Singapore (1998).
- [3] M. Cornacchia *et al.*, Linac Coherent Light Source (LCLS) Design Study Report, Report SLAC-R-521 (Stanford Linear Accelerator Center, Stanford, CA, revised 1998).
- [4] R. Brinkmann, G. Materlik, J. Rossbach, A. Wagner, Eds., *Conceptual Design of a 500 GeV  $e^+ e^-$  Linear Collider with Integrated X-Ray Laser Facility*, DESY Report DESY97-048 (Deutsches Elektronen-Synchrotron, Hamburg, 1997).
- [5] A. M. Kondratenko, E. L. Saldin, *Sov. Phys. Dokl.* **24** (no. 12), 986 (1979).
- [6] R. Bonifacio, C. Pellegrini, L. M. Narducci, *Opt. Commun.* **50**, 373 (1984).
- [7] J.B. Murphy, C. Pellegrini and R. Bonifacio, *Optics Comm.*: **53**, 197 (1985).
- [8] R. Bonifacio *et al.*, *Nucl. Instrum. Methods Phys Res. A* **293**, 627 (1990).
- [9] H. P. Freund *et al.*, *IEEE J. Quantum Electron.* **36**, 275 (2000).
- [10] Z. Huang, K. J. Kim, *Phys. Rev. E* **62**, 7295 (2000).
- [11] K.J. Kim and Z. Huang, "Present Status of X-ray FELs", to be published in the Proceeding of 19th ICFA Advanced Beam Dynamics Workshop on Future Light Source, Arcidosso, Italy, Sept. 10-15, 2000.
- [12] S. V. Milton *et al.*, *Phys. Rev. Lett.* **85**, 988 (2000).
- [13] R. Prazeres *et al.*, *Phys. Rev. Lett.* **78**, 2124 (1987).
- [14] M.J. Hogan, *et al.*, *Phys. Rev. Lett.* **80**, 289-292 (1998).
- [15] M.J. Hogan *et al.*, *Phys. Rev. Lett.* **81**, 4867 (1998).
- [16] D. C. Nguyen *et al.*, *Phys. Rev. Lett.* **81**, 810 (1998).
- [17] M. Babzien *et al.*, *Phys. Rev. E* **57**, 6093 (1998).
- [18] J. Andruszkow *et al.*, *Phys. Rev. Lett.* **85**, 3825 (2000).
- [19] S. Milton *et al.*, *Science*, **292** (2001) 2037-2041.
- [21] A. Tremaine *et al.*, in *Proceedings of the 2001 Particle Accelerator Conference*, Chicago, IL (2001).
- [22] A. Murokh *et al.*, in *Proceedings of the 2001 Particle Accelerator Conference*, Chicago, IL (2001).
- [23] V. Ayvazyan, *et al.*, submitted to *Nature*.
- [24] A. Tremaine *et al.*, *Phys. Rev. Lett.* **81**, 5816 (1998).
- [25] A. H. Lumpkin *et al.*, *Phys. Rev. Lett.* **86**, 79-82 (2001).

- [26] X.J Wang, X. Qiu, I. Ben-Zvi, Phys. Rev. E, **54**, R3121 – R3124 (1996).
- [27] R. Carr *et al.*, Phys. Rev. ST Accel. and Beams, **4**, 122402 (2001).
- [28] P. Emma *et al.*, in *Proceedings of the 20<sup>th</sup> International FEL Conference (FEL98)*, Williamsburg, Va., USA, 1198, SLAC-PB-7913.
- [29] A. Murokh *et al.*, in *Proceedings of the 22<sup>th</sup> International FEL Conference (FEL2000)*, Durham, NC, USA.
- [30] Z. Huang, Private Communication.
- [31] Z. Huang, K. J. Kim, Presented at 22<sup>nd</sup> International FEL conference (Durham, USA August 13-18 2000), to be published in Nucl. Instrum. Methods Phys Res. A.
- [32] S. Reiche, Nucl. Instr. & Meth. **A429**, 243(1999).
- [33] H.P. Freund, T.M. Antonsen Jr, Principle of Free-Electron Lasers (Chapman & Hall, London, ed. 2, 1996).
- [34] R. Kato *et al.*, *Proceedings of the 23<sup>rd</sup> International FEL Conference (FEL 2001)*, Darmstadt, Germany.




Figure 1(a). Single shot harmonic spectra of the fundamental, 2<sup>nd</sup> and 3<sup>rd</sup> harmonics




Figure 1(b). Harmonic Spectra independently measured at peak VISA gain.




Fig. 2. Energy vs. distance for the fundamental, 2<sup>nd</sup> and 3<sup>rd</sup> harmonics. The gain lengths for these are 19 cm, 9.8 cm, and 6.0 cm respectively.

