

Measurement of the Branching Fractions of $\Lambda_c^+ \rightarrow p\bar{K}n(\pi)$

CLEO Collaboration

Abstract

Using data recorded by the CLEO-II detector at CESR, we report new measurements of the branching fractions for the decays of the charmed baryon Λ_c^+ into $pK^-\pi^+\pi^0$, $p\bar{K}^0$, $p\bar{K}^0\pi^+\pi^-$, and $p\bar{K}^0\pi^0$, all measured relative to $pK^-\pi^+$. The relative branching fractions are $0.67 \pm 0.04 \pm 0.11$, $0.46 \pm 0.02 \pm 0.04$, $0.52 \pm 0.04 \pm 0.05$, and $0.66 \pm 0.05 \pm 0.07$ respectively.

M. S. Alam,¹ S. B. Athar,¹ Z. Ling,¹ A. H. Mahmood,¹ H. Severini,¹ S. Timm,¹
 F. Wappler,¹ A. Anastassov,² J. E. Duboscq,² D. Fujino,^{2,*} K. K. Gan,² T. Hart,²
 K. Honscheid,² H. Kagan,² R. Kass,² J. Lee,² M. B. Spencer,² M. Sung,² A. Undrus,^{2,†}
 R. Wanke,² A. Wolf,² M. M. Zoeller,² B. Nemati,³ S. J. Richichi,³ W. R. Ross,³ P. Skubic,³
 M. Bishai,⁴ J. Fast,⁴ J. W. Hinson,⁴ N. Menon,⁴ D. H. Miller,⁴ E. I. Shibata,⁴
 I. P. J. Shipsey,⁴ M. Yurko,⁴ L. Gibbons,⁵ S. Glenn,⁵ S. D. Johnson,⁵ Y. Kwon,^{5,‡}
 S. Roberts,⁵ E. H. Thorndike,⁵ C. P. Jessop,⁶ K. Lingel,⁶ H. Marsiske,⁶ M. L. Perl,⁶
 D. Ugolini,⁶ R. Wang,⁶ X. Zhou,⁶ T. E. Coan,⁷ V. Fadeyev,⁷ I. Korolkov,⁷ Y. Maravin,⁷
 I. Narsky,⁷ V. Shelkov,⁷ J. Staeck,⁷ R. Stroynowski,⁷ I. Volobouev,⁷ J. Ye,⁷ M. Artuso,⁸
 A. Efimov,⁸ M. Goldberg,⁸ D. He,⁸ S. Kopp,⁸ G. C. Moneti,⁸ R. Mountain,⁸ S. Schuh,⁸
 T. Skwarnicki,⁸ S. Stone,⁸ G. Viehhauser,⁸ X. Xing,⁸ J. Bartelt,⁹ S. E. Csorna,⁹ V. Jain,^{9,§}
 K. W. McLean,⁹ S. Marka,⁹ R. Godang,¹⁰ K. Kinoshita,¹⁰ I. C. Lai,¹⁰ P. Pomianowski,¹⁰
 S. Schrenk,¹⁰ G. Bonvicini,¹¹ D. Cinabro,¹¹ R. Greene,¹¹ L. P. Perera,¹¹ G. J. Zhou,¹¹
 B. Barish,¹² M. Chadha,¹² S. Chan,¹² G. Eigen,¹² J. S. Miller,¹² C. O'Grady,¹²
 M. Schmidtler,¹² J. Urheim,¹² A. J. Weinstein,¹² F. Würthwein,¹² D. W. Bliss,¹³
 G. Masek,¹³ H. P. Paar,¹³ S. Prell,¹³ V. Sharma,¹³ D. M. Asner,¹⁴ J. Gronberg,¹⁴
 T. S. Hill,¹⁴ D. J. Lange,¹⁴ S. Menary,¹⁴ R. J. Morrison,¹⁴ H. N. Nelson,¹⁴ T. K. Nelson,¹⁴
 C. Qiao,¹⁴ J. D. Richman,¹⁴ D. Roberts,¹⁴ A. Ryd,¹⁴ M. S. Witherell,¹⁴ R. Balest,¹⁵
 B. H. Behrens,¹⁵ W. T. Ford,¹⁵ H. Park,¹⁵ J. Roy,¹⁵ J. G. Smith,¹⁵ J. P. Alexander,¹⁶
 C. Bebek,¹⁶ B. E. Berger,¹⁶ K. Berkelman,¹⁶ K. Bloom,¹⁶ D. G. Cassel,¹⁶ H. A. Cho,¹⁶
 D. S. Crowcroft,¹⁶ M. Dickson,¹⁶ P. S. Drell,¹⁶ K. M. Ecklund,¹⁶ R. Ehrlich,¹⁶
 A. D. Foland,¹⁶ P. Gaidarev,¹⁶ B. Gittelmann,¹⁶ S. W. Gray,¹⁶ D. L. Hartill,¹⁶
 B. K. Heltsley,¹⁶ P. I. Hopman,¹⁶ J. Kandaswamy,¹⁶ P. C. Kim,¹⁶ D. L. Kreinick,¹⁶
 T. Lee,¹⁶ Y. Liu,¹⁶ G. S. Ludwig,¹⁶ N. B. Mistry,¹⁶ C. R. Ng,¹⁶ E. Nordberg,¹⁶ M. Ogg,^{16,**}
 J. R. Patterson,¹⁶ D. Peterson,¹⁶ D. Riley,¹⁶ A. Soffer,¹⁶ B. Valant-Spaight,¹⁶ C. Ward,¹⁶
 M. Athanas,¹⁷ P. Avery,¹⁷ C. D. Jones,¹⁷ M. Lohner,¹⁷ C. Prescott,¹⁷ J. Yelton,¹⁷
 J. Zheng,¹⁷ G. Brandenburg,¹⁸ R. A. Briere,¹⁸ A. Ershov,¹⁸ Y. S. Gao,¹⁸ D. Y.-J. Kim,¹⁸
 R. Wilson,¹⁸ H. Yamamoto,¹⁸ T. E. Browder,¹⁹ F. Li,¹⁹ Y. Li,¹⁹ J. L. Rodriguez,¹⁹
 T. Bergfeld,²⁰ B. I. Eisenstein,²⁰ J. Ernst,²⁰ G. E. Gladding,²⁰ G. D. Gollin,²⁰
 R. M. Hans,²⁰ E. Johnson,²⁰ I. Karliner,²⁰ M. A. Marsh,²⁰ M. Palmer,²⁰ M. Selen,²⁰
 J. J. Thaler,²⁰ K. W. Edwards,²¹ A. Bellerive,²² R. Janicek,²² D. B. MacFarlane,²²
 P. M. Patel,²² A. J. Sadoff,²³ R. Ammar,²⁴ P. Baringer,²⁴ A. Bean,²⁴ D. Besson,²⁴
 D. Coppage,²⁴ C. Darling,²⁴ R. Davis,²⁴ N. Hancock,²⁴ S. Kotov,²⁴ I. Kravchenko,²⁴
 N. Kwak,²⁴ S. Anderson,²⁵ Y. Kubota,²⁵ S. J. Lee,²⁵ J. J. O'Neill,²⁵ S. Patton,²⁵
 R. Poling,²⁵ T. Riehle,²⁵ V. Savinov,²⁵ and A. Smith²⁵

¹State University of New York at Albany, Albany, New York 12222

²Ohio State University, Columbus, Ohio 43210

³University of Oklahoma, Norman, Oklahoma 73019

⁴Purdue University, West Lafayette, Indiana 47907

*Permanent address: Lawrence Livermore National Laboratory, Livermore, CA 94551.

†Permanent address: BINP, RU-630090 Novosibirsk, Russia.

‡Permanent address: Yonsei University, Seoul 120-749, Korea.

§Permanent address: Brookhaven National Laboratory, Upton, NY 11973.

**Permanent address: University of Texas, Austin TX 78712

- ⁵University of Rochester, Rochester, New York 14627
- ⁶Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309
- ⁷Southern Methodist University, Dallas, Texas 75275
- ⁸Syracuse University, Syracuse, New York 13244
- ⁹Vanderbilt University, Nashville, Tennessee 37235
- ¹⁰Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
- ¹¹Wayne State University, Detroit, Michigan 48202
- ¹²California Institute of Technology, Pasadena, California 91125
- ¹³University of California, San Diego, La Jolla, California 92093
- ¹⁴University of California, Santa Barbara, California 93106
- ¹⁵University of Colorado, Boulder, Colorado 80309-0390
- ¹⁶Cornell University, Ithaca, New York 14853
- ¹⁷University of Florida, Gainesville, Florida 32611
- ¹⁸Harvard University, Cambridge, Massachusetts 02138
- ¹⁹University of Hawaii at Manoa, Honolulu, Hawaii 96822
- ²⁰University of Illinois, Champaign-Urbana, Illinois 61801
- ²¹Carleton University, Ottawa, Ontario, Canada K1S 5B6
and the Institute of Particle Physics, Canada
- ²²McGill University, Montréal, Québec, Canada H3A 2T8
and the Institute of Particle Physics, Canada
- ²³Ithaca College, Ithaca, New York 14850
- ²⁴University of Kansas, Lawrence, Kansas 66045
- ²⁵University of Minnesota, Minneapolis, Minnesota 55455

Since the first observation of the lowest lying charmed baryon, the Λ_c^+ , there have been many measurements made of its exclusive decay channels. As it is difficult to measure the production cross-section of the Λ_c^+ baryons, decay rates are typically presented as branching ratios relative to $\Lambda_c^+ \rightarrow pK^-\pi^+$, the most easily observed decay channel. However, fewer than half of the Λ_c^+ hadronic decays are presently accounted for. Measurement of these modes is of practical as well as theoretical interest. Here, we present measurements of the branching fractions of Λ_c^+ into $pK^-\pi^+\pi^0$, $p\bar{K}^0$, $p\bar{K}^0\pi^+\pi^-$, and $p\bar{K}^0\pi^0$, all relative to $pK^-\pi^+$. The last of these is the first measurement of this mode. The other modes have been previously measured but with considerably larger uncertainties than in the present study.

The data presented here were taken by the CLEO II detector [1] operating at the Cornell Electron Storage Ring. The sample used in this analysis corresponds to an integrated luminosity of $4.8 fb^{-1}$ from data taken on the $\Upsilon(4S)$ resonance and in the continuum at energies just above and below the $\Upsilon(4S)$. We detected charged tracks with a cylindrical drift chamber system inside a solenoidal magnet. Photons were detected using an electromagnetic calorimeter consisting of 7800 cesium iodide crystals.

Particle identification of p , K^- , and π^+ candidates was performed using specific ionization measurements in the drift chamber, and when present, time-of-flight measurements. For each mass hypothesis, a combined χ^2 probability P_i was formed ($i = \pi, K, p$). Using these probabilities, a normalized probability ratio L_i was evaluated, where $L_i = P_i/(P_\pi + P_K + P_p)$. Well identified protons peaked near $P_p = 1.0$ while tracks that were identified to not be protons peak near $P_p = 0.0$. For a track to be used as a proton in this study, we required it to have $L_p > 0.8$, which eliminated much of the background, though with considerable diminution of efficiency. For kaons we applied a looser and more efficient cut of $L_K > 0.1$. We have chosen these cuts using a Monte Carlo simulation program to maximize the significance of the signals. The proton identification requirement resulted in an efficiency that is strongly momentum dependent, whereas the kaon identification is rather efficient at all momenta. In order to reduce the large combinatoric background, we required $x_p > 0.5$, where $x_p = P_{\Lambda_c}/\sqrt{E_{beam}^2 - m_{\Lambda_c}^2}$ is the scaled momentum of the Λ_c^+ candidate. Approximately 60% of Λ_c^+ baryons from $c\bar{c}$ continuum events passed this requirement.

The \bar{K}^0 candidates were identified in their decay $K_s^0 \rightarrow \pi^+\pi^-$, by reconstructing a secondary vertex from the intersection of two oppositely charged tracks in the $r - \phi$ plane. The invariant mass of the \bar{K}^0 candidate must lie within 9 MeV/ c^2 (around 3 standard deviations (σ) of its nominal value.

The π^0 candidates were selected through their decay $\pi^0 \rightarrow \gamma\gamma$ from pairs of well-defined showers in the CsI calorimeter with a reconstructed invariant mass within 3σ of the π^0 mass. In order to reduce the combinatorial background, each γ was required to have an energy of at least 50 MeV, and the π^0 was required to have a momentum of at least 300 MeV/ c .

The resulting mass distributions for the 5 modes are shown in Figure 1. Each peak was fit to the sum of a Gaussian signal distribution with width fixed to that obtained from CLEO's GEANT based Monte Carlo simulation program, and a second order polynomial background distribution. The signal widths used and the resulting signal yields are tabulated in Table 1.

The efficiency for each Λ_c^+ mode was calculated using the Monte Carlo simulation program [2]. The particle identification efficiency was checked using real data from $\Lambda \rightarrow p\pi^-$ and

Mode	MC Width (MeV)	Signal
$pK^-\pi^+$	16	10109 ± 191
$pK^-\pi^+\pi^0$	22	2606 ± 165
$p\bar{K}^0$	19	1025 ± 40
$p\bar{K}^0\pi^+\pi^-$	15	985 ± 65
$p\bar{K}^0\pi^0$	27	774 ± 52

TABLE I. The number of Λ_c^+ 's found with $x_p(\Lambda_c) > 0.5$

$D^{*+} \rightarrow K^-\pi^+\pi^+$ decays that were identified topologically. The reconstruction efficiency of the Λ_c^+ decays has some dependence on the resonant substructure of these states. In the case of the $pK^-\pi^+$ mode, the Monte-Carlo generator produced a mixture of non-resonant three-body decay together with $\Delta^{++}K^-$ and $p\bar{K}^{*0}$ decays, according to their measured branching fractions [3]. These three types of decays had slightly different reconstruction efficiencies, so that including the substructure changes the efficiency by $\Delta\epsilon/\epsilon = 0.02$ relative to 3-body phase space. We have also investigated the dependence of the reconstruction efficiency of the other modes on possible resonant substructure. The poor signal to background ratio did not allow a detailed measurement of the substructure of these modes. The efficiency calculation took into account the $\bar{K}^0 \rightarrow K_s^0$ and $K_s^0 \rightarrow \pi^+\pi^-$ branching fractions.

We have considered many possible sources of systematic error in the measurement. The main contributors to the systematic uncertainty came from the following sources: 1) Uncertainties in the fitting procedures, which were estimated by looking at the changes in the yields using different orders of polynomial background and different signal widths (15% in the case of $pK^-\pi^+\pi^0$, but much smaller for the other modes), 2) uncertainties due to the unknown mix of resonant substructure in the multi-body decays (up to 3% depending on the mode), 3) uncertainties due to π^0 finding (5%), K_s^0 finding (5%) and track finding (1%), and 4) uncertainties in the reconstruction efficiency due to the particle identification criteria for protons and kaons (4%). These uncertainties have been added in quadrature to obtain the total systematic uncertainty for each mode, taking into account the fact that many of these tend to cancel in a measurement of ratios of branching fractions.

There are three main types of quark decay diagrams that contribute to Λ_c^+ decays. The simplest method is the simple spectator diagram in which the virtual W^+ fragments independently of the spectator quark. The second method involves the quark daughters of the W^+ combining with the remaining quarks. The third method, W-exchange, involves the W^+ combining with the initial d quark. Unfortunately all the decay modes under investigation here can proceed by more than one of these decay diagrams, and their decay rates are not amenable to calculation.

In conclusion, we have measured new branching fractions of the Λ_c^+ into 4 decay modes, measured relative to the normalizing mode $\Lambda_c^+ \rightarrow pK^-\pi^+$. The results for three of these modes are in agreement with, and more accurate than, previous measurements. We have made the first measurement of the decay rate of $\Lambda_c^+ \rightarrow p\bar{K}^0\pi^0$. These measurements help account for the total width of the Λ_c^+ and increase the understanding of charmed baryon decays.

Mode	Relative Efficiency	$B/B(pK^-\pi^+)$	Previous Measurements
$pK^-\pi^+$	1.0	1.0	
$pK^-\pi^+\pi^0$	0.383	$0.67 \pm 0.04 \pm 0.11$	$0.72^{+0.32}_{-0.22}$ [5]
$p\bar{K}^0$	0.218	$0.46 \pm 0.02 \pm 0.04$	$0.44 \pm 0.07 \pm 0.05$ [4] $0.55 \pm 0.17 \pm 0.14$ [6] $0.62 \pm 0.15 \pm 0.03$ [7]
$p\bar{K}^0\pi^+\pi^-$	0.187	$0.52 \pm 0.04 \pm 0.05$	$0.43 \pm 0.12 \pm 0.04$ [4] $0.98 \pm 0.36 \pm 0.08$ [5]
$p\bar{K}^0\pi^0$	0.115	$0.66 \pm 0.05 \pm 0.07$	

TABLE II. The measured relative branching fractions

REFERENCES

- [1] Y. Kubota et al., Nucl. Inst. and Methods A320, 66 (1992).
- [2] R. Brun et al., CERN/DD/EE/84-11.
- [3] Review of Particle Properties, R. Barnett et al., Phys. Rev. D54 1 (1996).
- [4] P. Avery et al., Phys. Rev. D43, 3499 (1991)
- [5] S. Barlag et al., Z. Phys. C48, 29 (1990).
- [6] J. Anjos et al., Phys. Rev. D41, 801 (1990).
- [7] H. Albrecht et al., Phys. Lett. B207, 109 (1988).




FIG. 1. Invariant mass plots for the 5 different decay modes of the Λ_c^+