Sensitivity to New Physics in the W + jet-jet Decay Channel at the Tevatron

Dave Toback

University of Chicago

ABSTRACT

Anomalous production of W + jet-jet events is possible at the Tevatron in many theories of new physics such as technicolor and extended gauge models. We outline methods for searching for new heavy particles, X, via the decay $X \rightarrow Wjj$ with and without jet b-tagging. We estimate the sensitivity to $W' \rightarrow WZ$, $Z' \rightarrow WW$ and $\rho_T \rightarrow Wjj$ for Run II (2 fb⁻¹) and TeV33 (30 fb⁻¹) and discuss some of the challenging experimental problems associated with searching for technicolor signatures.

I. INTRODUCTION

W', Z' and technicolor models [1, 2] have long been known to predict new heavy particles which in turn decay into the W+jetjet final state at the Tevatron [3]. The W' search in the Run 1 data is complete, so we use it to outline the search methods and, using some simple assumptions, we extrapolate the sensitivity to 30 fb⁻¹ of data. For those cases which have final state b-quarks, we discuss possibilities using b-tagging. Finally, we discuss some of the challenging problems associated with searching for technicolor signatures.

II. THEORY

A. Heavy Vector Bosons W' and Z'

In the simplest W' model, known as the reference model [1], the W' is a heavier version of the ordinary W and is assumed to have the same vertex couplings ($Wq\bar{q}, Wl\nu$ and WWZ) as the ordinary W in the Standard Model (SM). Also all fermions, specifically neutrinos, are the same as their SM counterparts. The dominant features of this model are the high production cross sections and the increase in $\Gamma(W' \to WZ)$ as $M_{W'}^5$ which gives rise to a large branching fraction into WZ. However, at large masses ($M_{W'} \approx 425$ GeV) the resonance width becomes so wide ($\Gamma \approx \frac{M_{W'}}{2}$) that perturbation theory breaks down[4] and the model is no longer applicable. This model has already been excluded in the Run 1 data [5].

In extended gauge models[1], proposed to restore left-right symmetry to the weak force, the effective W'WZ vertex becomes parameterized in terms of a mixing angle ξ which is estimated to be $\xi = C(\frac{M_W}{M_{W'}})^2$ where *C* is a constant of order 1[6]. In this case the width only increases linearly with $M_{W'}$. In this note all results assume *C*=1. Similar arguments hold for the production and decay of $Z' \to WW \to e\nu jj$. In the mass region above 200 GeV, as shown in [1], $\sigma \cdot Br(Z' \to e\nu jj) \approx$ $\sigma \cdot Br(W' \to e\nu jj)$.

B. Technicolor

We assume a simple toy model of color singlet technicolor[2], which has a technirho ($\rho_{\rm T}$) which decays into a pair of technipions ($\Pi_{\rm T}$). In this model, there is an isotriplet of mass-eigenstate technipions $\pi_{\rm T}^{\pm}$ and $\pi_{\rm T}^{0}$ which mix with $W_{\rm L}^{\pm}$ and $Z_{\rm L}^{0}$ via $|\Pi_{\rm T}\rangle =$ $\sin \chi |W_{\rm L}\rangle + \cos \chi |\pi_{\rm T}\rangle$, where χ is a mixing angle. Thus, depending on the masses and the value of the mixing angle, the following decay modes are allowed:

Charged decays

$$\begin{array}{l} \rho_{\rm T}^+ \rightarrow W^+ Z^0 \leftarrow \\ \rho_{\rm T}^+ \rightarrow W^+ \pi_{\rm T}^0 \leftarrow \\ \rho_{\rm T}^+ \rightarrow Z^0 \pi_{\rm T}^+ \\ \rho_{\rm T}^+ \rightarrow \pi_{\rm T}^0 \pi_{\rm T}^+ \end{array}$$

- Neutral decays
 - $\begin{array}{c} \rho_{\rm T}^{\rm 0} \rightarrow W^+ W^- \leftarrow \\ \rho_{\rm T}^{\rm 0} \rightarrow W^+ \pi_{\rm T}^- \leftarrow \\ \rho_{\rm T}^{\rm 0} \rightarrow W^- \pi_{\rm T}^+ \leftarrow \\ \rho_{\rm T}^{\rm 0} \rightarrow \pi_{\rm T}^+ \pi_{\rm T}^- \end{array}$

where the arrows indicate the channels which have possible W_{jj} final states.

Technipion decays are Higgs-like, i.e, the dominant decay is expected to be into heavy fermion pairs:

- $\pi_{\rm T}^0 \rightarrow b\bar{b}$ if $M_{\pi_{\rm T}} < 2m_t$
- $\pi_{\mathrm{T}}^{0} \rightarrow t\bar{t}$ if $M_{\pi_{\mathrm{T}}} > 2m_{t}$
- $\pi_{\rm T}^+ \rightarrow c \bar{b}, c \bar{s} \text{ or } \tau \nu \text{ if } {\rm M}_{\pi_{\rm T}} < m_t + m_b$
- $\pi^+_{
 m T}
 ightarrow t ar{b}$ if ${
 m M}_{\pi_{
 m T}} > m_t + m_b$

These processes can be broken down into three final state jj cases: 2 b-quarks, 1 b-quark and 0 b-quarks. However, the cross sections and branching ratios are HIGHLY dependent on the masses of the $\rho_{\rm T}$ and the $\pi_{\rm T}$ as well as the mixing angle between the vector bosons and the mass-eigenstate $\pi_{\rm T}$'s.

There are many challenges associated with looking for technicolor signatures. There are (at least) three unknown parameters in the theory; M_{ρ} , M_{π} , and χ , which make it unclear where it is we should look. However, many of the final state signatures are similar and can be searched for simultaneously. Another problem with multiple similar signatures is that for some combination of the parameters, many of the final states would look similar and perhaps even completely wash out a signal. For example, if $M_{\rho} = 400$ GeV, $M_{\pi} = 110$ GeV, and the mixing angle is such that $\sigma \cdot Br(\rho \rightarrow WW) \approx \sigma \cdot Br(\rho \rightarrow WZ) \approx \sigma \cdot Br(\rho \rightarrow W\pi)$ we could have three resonances so close that we might misinterpret the events as a continuum and part of a small normalization problem.

III. WHAT'S BEEN DONE SO FAR: THE SEARCH FOR $W' \rightarrow WZ$

The Run 1 $W' \rightarrow WZ$ search is complete[5], so we use it to outline the general W + jet-jet search, and making some simple assumptions, extrapolate the sensitivity to 30 fb⁻¹ of data.

We highlight the main features of the methods used in the $W' \rightarrow WZ$ search and point out the primary limitations to using this as an example for technicolor:

- The search identifies the W via the decay channel $W \rightarrow e\nu$;
- The search identifies the Z via the decay[7] channel $Z \rightarrow jj$
- It uses high kinematic thresholds to reduce the W+Jets backgrounds
- The limits already set assume that the new production would show up as ONE bump in the jet-jet mass spectrum (coming from a Z boson and centered at 90 GeV) vs. W+jet-jet mass[8] spectrum (coming from a new particle and centered at its mass)
- When the total widths are well defined and small, the limits used in the extended gauge model are applicable
- The method was designed for good acceptance for the total event mass in the region around 500 GeV. It is not clear that this is the optimal method for lower mass configurations
- The method is very general and can be used to search for any high-mass jet-jet vs. W+jet-jet mass spectral anomaly (i.e, from any new particle or particles X, Y, which might decay via X → WY → Wjj)
- The region 'outside the peak' is used to normalize the W+Jets cross section
- Using these cuts, roughly 20% of the events are from backgrounds other than SM W+Jets

We summarize the cuts here:

- electron $E_T \ge 30$ GeV, $|\eta| < 1.0$
- At least two high energy jets. (1 Jet with $E_T \ge 50$ GeV, a second Jet with $E_T \ge 20$ GeV. Both with $|\eta| < 2.0$)

IV. RESULTS

A. Looking at the data in Run 1

Figure 1 shows the W+jet-jet mass spectra for the data and background from Run 1. We note that the number of W+jets events is normalized to the number of data events minus the other backgrounds. The data are well modeled by the background Monte Carlos. If there were new particle production $(W', Z' \text{ or } \rho_T)$ with a large enough cross section, it would show up as a bump in this distribution.




Figure 1: W+jet-jet mass spectrum for the data along with the background predictions. The background spectrum is normalized to the same number of events in the data as in the backgrounds. There is no evidence for W' or any new particle produced in association with a W. The $\chi^2/d.o.f. = 1.30$. We expect 4.2 events above $M_{W+jj} > 600$ GeV; we observe 7.


B. Results from the $W' \rightarrow WZ$ Search

The acceptance rises from 6% at $M_{W'} = 200 \text{ GeV/c}^2$ to 35% at $M_{W'} = 600 \text{ GeV/c}^2$, and the 95% CL exclusion limits in the mass vs. ξ plane are shown in Figure 2. Since we have excluded the reference model, we concentrate on the sensitivity to the extended gauge model with $\xi = (\frac{M_W}{M_{W'}})^2$. For simplicity, because this is a background limited exclusion, we have made the reasonable assumption that the cross section limits scale as the inverse of the square root of the luminosity.

Similar arguments hold for the production and decay of $Z' \rightarrow WW \rightarrow e\nu jj$. Since the W and the Z have similar masses we expect that the limits are virtually identical to those of the W', i.e, we assume $\sigma_{95\%CL} \cdot Br(Z' \rightarrow e\nu jj) = \sigma_{95\%CL} \cdot Br(W' \rightarrow e\nu jj)$. These results are shown in Figure 3.

C. Extrapolating Results to the $\rho_{\rm T} \rightarrow WX \rightarrow e\nu jj$ Search

As previously mentioned, the masses as well as the branching ratios to the various decay modes are highly model dependent. For simplicity, we assume that the limits for $\rho_T \rightarrow WX \rightarrow e\nu jj$ are identical to that for the W' case and that they scale






Figure 2: 95% CL upper limit of $\sigma \cdot Br(W' \to WZ) \cdot Br(W \to e\nu)$ vs. $M_{W'}$ for the extended gauge model with $\xi = (\frac{M_W}{M_{W'}})^2$. The solid line is the theoretically expected $\sigma \cdot Br$. The dashed lines show predicted limits for 110 pb⁻¹, 2 fb⁻¹ and 30 fb⁻¹ respectively. Note that we have assumed that the limits simply scale as the inverse of the square root of the luminosity.

as the inverse of the square root of the luminosity. For the theoretical predictions we assume that the branching ratio $\rho_T \rightarrow WX \rightarrow Wjj = 100\%$, and that the mass of the intermediate particle, X, is near that of the Z. These results are shown in Figure 4.

V. WHEN THERE ARE FINAL STATE B-QUARKS

A. Overview

The $\rho_{\rm T}$ often decays to $\pi_{\rm T}$ which in turn can decay to bquarks. The $e\nu bj$ and $e\nu bb$ final states are very different from the generic $e\nu jj$ final states as they have the additional final state b-quark which can be tagged. Since there could be either one or two final state b's we use an event selection similar to that of the CDF top analysis [9]. This selection criteria has a number of advantages and disadvantages:

Advantages

Less background;

We can an lower kinematic thresholds to get more acceptance;

Figure 3: 95% CL upper limit of $\sigma \cdot Br(Z' \to WW) \cdot Br(W \to e\nu)$ vs. $M_{Z'}$ for the extended gauge model with $\xi = (\frac{M_Z}{M_{Z'}})^2$. The solid line is the theoretically expected $\sigma \cdot Br$. The dashed lines show predicted limits for 110 pb⁻¹, 2 fb⁻¹ and 30 fb⁻¹ respectively. We have assumed $\sigma_{95\% CL} \cdot Br(Z' \to e\nu jj) = \sigma_{95\% CL} \cdot Br(W' \to e\nu jj)$ and that the limits simply scale as the inverse of the square root of the luminosity.

Cleaner signal

• Disadvantages

Smaller Branching Ratio into final state b-quarks.

Smaller Efficiency Acceptance

Looser Kinematic thresholds require tighter lepton identification cuts.

We use the following cuts:

- Central, isolated electron or muon ($|\eta| < 1.0$)
- Tight electron and Muon identification cuts [9]
- Lepton $E_T > 20 \text{ GeV}$
- ₽_T > 20 GeV
- One or more b-tagged jet




Figure 4: 95% CL upper limit of $\sigma \cdot Br(\rho_T \to Wjj)$ vs. M_{ρ_T} . The solid line is the theoretically expected $\sigma \cdot Br$ for the color singlet case and assume $\rho_T \to WX \to Wjj = 100\%$. The dashed lines show predicted limits for 110 pb⁻¹, 2 fb⁻¹ and 30 fb⁻¹ respectively. Note that we have assumed that the limits simply scale as the inverse of the square root of the luminosity

B. Limit Estimates

We expect roughly 50 events passing the above cuts for Run 1, and the efficiency × Acceptance for these cuts is $\approx 0.5\%$ for $\pi_{\rm T} \rightarrow b\bar{b}$ with $M_{\pi_{\rm T}} = 100$ GeV and $M_{\rho_{\rm T}} = 200$ GeV. To set limits we use:

$$\sigma = rac{\mathrm{N}_{\mathrm{Events}}}{\mathrm{Lum}\cdot\mathrm{Acc}\cdot\epsilon}$$

Two methods are used to estimate the limits:

• Assume all are 'signal' to set limits (Conservative):

$$95\%$$
 CL Upper Event Limit = 64 events

$$\rightarrow \sigma \cdot Br(\rho_{\rm T} \rightarrow l\nu bb) = 120 \text{ pb.}$$

• Assume all are 'background' to set limits:

$$\rightarrow \sigma \cdot Br(\rho_{\rm T} \rightarrow l\nu bb) = 30 \text{ pb.}$$

Scaling by the square root of the luminosity we find limits of 2 pb with 30 fb^{-1} . However, we could probably do better with fits and smarter cuts.

VI. CONCLUSIONS

We have outlined methods for searching for new heavy particles, X, via the decay $X \rightarrow Wjj$ and presented a summary of the sensitivity to new physics for luminosities up to 30 fb⁻¹ for heavy vector bosons (W' and Z') and for color singlet technicolor. Since some cases have final state b-quarks, we presented some preliminary methods and results using b-tagging. In addition we have highlighted some of the problems associated with searching for technicolor signatures. The bottom line is that the W+jet-jet channel is a good place to look for new physics. In fact there are so many things which could show up that we wouldn't even know what we had if we did find something. If there is nothing to be found we will have sensitivity to these processes at the 300 - 500 GeV level with 30 fb⁻¹ of data.

VII. ACKNOWLEDGMENTS

The author would like to thank Henry Frisch, Sacha Kopp, Jose Benlloch, Rob Harris, Tom Rizzo, JoAnne Hewett and Ken Lane for their help in this analysis.

VIII. REFERENCES

- G. Altarelli, B. Mele and M. Ruiz-Altaba Searching for new Heavy Vector Bosons in PP colliders Z. Phys. C45, P109-P121 (1989)
- [2] E. Eichten and K. Lane *Two Scale Technicolor Physics Letters* 99B (1980). K. Lane and M.V. Ramana *Walking Technicolor* signatures at Hadron Colliders Phys. Rev. D 44, 2678 (1991) E. Eichten and K. Lane Low-Scale Technicolor at the Tevatron FERMILAB-PUB-96/075-T, BUHEP-96-9, hep-ph/9607213; to appear in Physics Letters B. E. Eichten and K. Lane Electroweak and Flavor Dynamics at Hadron Colliders–I FERMILAB-PUB-96/297-T, BUHEP-96-33, hep-ph/9609297; to appear in the proceedings of the 1996 DPF/DPB Summer Study on New Directions for High Energy Physics (Snowmass 96).
- [3] We note that these methods are applicable for searches such as $W' \rightarrow tb, W' \rightarrow WH, H \rightarrow WW$ and associated Higgs (*WH*) production.
- [4] J. Gunion, H. Haber, G. Kane and S. Dawson, *The Higgs Hunter's Guide*. Frontiers in Physics series. 1990.
- [5] T. Kamon (CDF Collaboration), Fermilab-Conf-96/106-E, Proceedings of the XXXIst Rencontres de Moriond (QCD), Les Arcs, Savoie, France, March 23-30, 1996.
- [6] P. Ramond, Ann. Rev. Nucl. Part. Sci. 33 (1983) 31.
- [7] Note that we are not requiring the jets to be hadronic decays from quarks. Specifically, Z⁰ → e⁺e⁻ and Z⁰ → τ⁺τ⁻ decays are included in this search by virtue of the fact that we are simply searching for two energy clusters in the detector.

of the electron-neutrino system to be equal to W boson mass[10] (taken to be 80 GeV/c²) then P_z^{ν} is restricted to two possible values. When the transverse mass is greater than 80 GeV/c², the two an P_z^{ν} solutions have imaginary components. In such events the constraint is made to the value of the transverse mass instead of to 80 GeV/c² which gives two identical solutions. Once we have the neutrino solution(s), we find the invariant mass of the W+jet-jet system by using the two jets. We choose the solution with the lower invariant mass since studies have shown that it does a better job of giving back the correct W' mass and width. We also use the same procedure on the data as for the background and the signal Monte Carlo samples to reduce any other systematic problems introduced by this method.

- [9] F. Abe et al., Phys. Rev. D 50, 2966 (1994)
- [10] F. Abe et al., Phys. Rev. Lett. 73 2296 (1994)